



November 9, 2001 / Vol. 50 / No. RR-19



***Recommendations
and
Reports***

Inside: Two Continuing Education Examinations

Revised Guidelines for HIV Counseling, Testing, and Referral

and

Revised Recommendations for HIV Screening of Pregnant Women

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention (CDC)
Atlanta, GA 30333



All *MMWR* references are available on the Internet at <<http://www.cdc.gov/mmwr/>>. Use the search function to find specific articles.

Single copies of this document are available free of charge from the HIV/AIDS Treatment Information Service (ATIS) and can be obtained by calling (800) 448-0440, (301) 519-0459 (international), or (888) 480-3739 (TTY) or by downloading the document from the ATIS website at <www.hivatis.org>.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites.

MMWR

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy on Friday of each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/mmwr/> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/Publications/mmwr/>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to: Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone (888) 232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

The *MMWR* series of publications is published by the Epidemiology Program Office, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

Centers for Disease Control and Prevention. Revised Guidelines for HIV Counseling, Testing, and Referral and Revised Recommendations for HIV Screening of Pregnant Women. *MMWR* 2001;50(No. RR-19):[inclusive page numbers].

Centers for Disease Control and Prevention Jeffrey P. Koplan, M.D., M.P.H.
Director

The material in this report was prepared for publication by
National Center for HIV, STD, and TB Prevention Harold W. Jaffe, M.D.
Acting Director

Division of HIV/AIDS Prevention — Surveillance
and Epidemiology Robert S. Janssen, M.D.
Director

The production of this report as an *MMWR* serial publication was coordinated in
Epidemiology Program Office Barbara R. Holloway, M.P.H.
Acting Director

Office of Scientific and Health Communications John W. Ward, M.D.
Director
Editor, MMWR Series

Recommendations and Reports Suzanne M. Hewitt, M.P.A.
Managing Editor

Amanda Crowell
Project Editor

Lynda G. Cupell
Visual Information Specialist

Michele D. Renshaw
Erica R. Shaver
Information Technology Specialists

Contents

Revised Guidelines for HIV Counseling, Testing, and Referral	1
Introduction	1
Targeted Versus Routinely Recommended HIV CTR	7
HIV Counseling	13
HIV Testing	27
HIV Referral	36
HIV CTR Services in Nontraditional Settings	39
Quality Assurance and Evaluation of HIV CTR Services	43
Conclusion	45
Additional Resources	45
References	46
Glossary	54
Continuing Education Examination	CE-1-19a1
Revised Recommendations for HIV Screening of Pregnant Women	59
Introduction	63
Background	65
Recommendations	75
Conclusions	81
References	81
Continuing Education Examination	CE-1-19a2

**Technical Expert Panel Review of CDC
HIV Counseling, Testing, and Referral Guidelines
February 18–19, 1999
Atlanta, Georgia**

Terje J. Anderson
National Association of People with AIDS
Washington, D.C.

David Atkins, M.D., M.P.H.
Agency for Healthcare Research and Quality
Rockville, Maryland

Catherine Baker-Cirac
California Office of AIDS
Sacramento, California

Ronald Bayer, Ph.D.
Columbia University
New York, New York

Frank K. Beadle de Palomo, M.A.
Academy for Educational Development
Washington, D.C.

Gail A. Bolan, M.D.
California Department of Health
Berkeley, California

Carol A. Browning, M.S.
Rhode Island Department of Health
Providence, Rhode Island

Scott Burris, J.D.
Temple University
Philadelphia, Pennsylvania

Amy S. DeGross, M.P.H.
Centers for Disease Control and Prevention
(CDC)
Atlanta, Georgia

John M. Douglas, M.D.
Denver Public Health
Denver, Colorado

Martin Fishbein, Ph.D.
University of Pennsylvania
Philadelphia, Pennsylvania

Alice A. Gandelman, M.P.H.
California STD Control Branch
Berkeley, California

Cynthia A. Getty
CDC
Atlanta, Georgia

Lawrence O. Gostin, J.D., L.L.D.
Georgetown University
Washington, D.C.

Khurram S. Hassan, M.P.H.
United Way of Metro Atlanta
Atlanta, Georgia

Thomas L. Hearn, M.S., Ph.D.
CDC
Atlanta, Georgia

Michael P. Johnson, M.D., M.P.H.
Health Resources and Services
Administration
Rockville, Maryland

William J. Kassler, M.D., M.P.H.
New Hampshire Department of Health
& Human Services
Concord, New Hampshire

Marlene LaLota, M.P.H.
Florida Department of Health
Tallahassee, Florida

Michael K. Lindsay, M.D., M.P.H.
Emory University
Atlanta, Georgia

Michael H. Merson, M.D.
Yale University
New Haven, Connecticut

Stephen F. Morin, M.A., Ph.D.
University of California, San Francisco
San Francisco, California

James Pearson, M.P.H., Ph.D.
Division of Consolidated Laboratory
Services
Richmond, Virginia

Beny J. Primm, M.D.
Urban Resources Institute
Brooklyn, New York

Joel Rosenstock, M.D., M.P.H.
Infectious Disease Solutions, P.C.
Atlanta, Georgia

Peter Salovey, Ph.D.
Yale University
New Haven, Connecticut

Charles A. Schable, M.S.
CDC
Atlanta, Georgia

Kathleen J. Sikkema, Ph.D.
Yale University
New Haven, Connecticut

Edith Springer, M.S.W.
Edith Springer Associates
Brooklyn, New York

Janis Spurlock-McLendon, M.S.W.
Connecticut Department of Public Health
Hartford, Connecticut

Lee Trevithick, M.A.
Cocoon House
Everett, Washington

James Welch
Division of Public Health
Dover, Delaware

The following CDC staff members prepared this report:

Beatrice T. Divine, M.A.

Stacie M. Greby, D.V.M., M.P.H.

Kenneth V. Hunt

Mary L. Kamb, M.D., M.P.H.

Richard W. Steketee, M.D., M.P.H.

Lee Warner, M.P.H.

*Division of HIV/AIDS Prevention — Surveillance and Epidemiology
National Center for HIV, STD, and TB Prevention***in consultation with**

Liisa M. Randall, M.A.

National Alliance of State and Territorial AIDS Directors

Revised Guidelines for HIV Counseling, Testing, and Referral

Summary

These guidelines replace CDC's 1994 guidelines, HIV Counseling, Testing, and Referral Standards and Guidelines, and contain recommendations for public- and private-sector policy makers and service providers of human immunodeficiency virus (HIV) counseling, testing, and referral (CTR). To develop these guidelines, CDC used an evidence-based approach advocated by the U.S. Preventive Services Task Force and public health practice guidelines. The recommendations are based on evidence from all available scientific sources; where evidence is lacking, opinion of "best practices" by specialists in the field has been used.

This revision was prompted by scientific and programmatic advances in HIV CTR, as well as advances in prevention and the treatment and care of HIV-infected persons. These advances include a) demonstrated efficacy of HIV prevention counseling models aimed at behavioral risk reduction; b) effective treatments for HIV infection and opportunistic infections; c) effective treatment regimens for preventing perinatal transmission; and d) new test technologies.

Although the new guidelines include many aspects of the previous ones (e.g., encouragement of confidential and anonymous voluntary HIV testing, need for informed consent, and provision of HIV prevention counseling that focuses on the client's own risk), the new guidelines differ in several respects, including

- giving guidance to all providers of voluntary HIV CTR in the public and private sectors;*
- using an evidence-based approach to provide specific recommendations for CTR;*
- underscoring the importance of early knowledge of HIV status and making testing more accessible and available;*
- acknowledging providers' need for flexibility in implementing the guidelines, given their particular client base, setting HIV prevalence level, and available resources;*
- recommending that CTR be targeted efficiently through risk screening and other strategies; and*
- addressing ways to improve the quality and provision of HIV CTR.*

INTRODUCTION

Purpose of the Guidelines

These guidelines were developed for policy makers and service providers in the many settings that offer voluntary human immunodeficiency virus (HIV) counseling, testing, and referral (CTR) — public and private, urban and rural, and those with high and low HIV prevalence (Box 1). The guidelines are intended to be used to develop CTR services and policies in traditional clinical settings (e.g., sexually transmitted disease

BOX 1. HIV counseling, testing, and referral (CTR) settings

Settings that provide HIV CTR include but are not limited to the following traditional clinical and nontraditional settings:

- Adolescent health clinics, school-based health centers, university health centers
- AIDS services organizations
- Clinics serving men who have sex with men
- Community-based organizations
- Community health centers
- Correctional facilities
- Drug or alcohol prevention and treatment programs
- Family planning clinics
- Freestanding HIV test sites
- Hospital emergency departments
- Hospitals/other urgent care centers
- Managed care organizations
- Men's health clinics
- Migrant health centers
- Occupational/employee health clinics
- Outreach programs (e.g., syringe exchange programs)
- Prenatal clinics
- Private-sector service providers
- Publicly funded counseling and testing sites
- Sexually transmitted disease clinics
- Tuberculosis clinics
- Women's health clinics

[STD] clinics, private physicians' offices) and nontraditional settings (e.g., community-based or outreach settings [homeless shelters, bars]), which can be important places to provide access to CTR to persons at increased HIV risk. The Public Health Service is responsible for ensuring the quality of services in publicly funded programs, and many aspects of these guidelines focus on such programs. The guidelines could also be useful for CTR in other settings (e.g., for insurance, military, and blood donation purposes). Recommendations should be tailored to fit the needs of clients, communities, and programs within local, state, and federal rules and regulations.

Evolution of the Guidelines

These guidelines revise and update several sets of CDC guidelines for HIV CTR. The first CDC guidelines, published in 1986, highlighted the importance of offering voluntary testing and counseling and maintaining confidential records (1). In 1987, new guidelines emphasized the need to decrease barriers to counseling and testing, especially disclosure of personal information (2). In 1993, an additional report described the model of HIV prevention counseling currently recommended — an interactive rather than didactic model focusing on a personalized HIV risk-reduction plan (3). In 1994, *HIV Counseling, Testing and Referral Standards and Guidelines* focused on standard counseling and testing procedures and reiterated the importance of the HIV prevention counseling model and the need for confidentiality of counseling (4).

Because of recent advances in HIV treatment and prevention (5–32, *Revised Recommendations for HIV Screening of Pregnant Women*), CDC consulted with multiple partners to revise the 1994 guidelines using an evidence-based approach (33,34) and to expand the target audience to all providers of HIV CTR in the United States (33). Where scientific findings were lacking, recommendations were guided by “best practices” from specialists in the field. These guidelines were developed through the following five-step approach:

- **Address user needs.** A survey was conducted of publicly funded sites that offer HIV CTR to assess user satisfaction with the 1994 CDC guidelines for HIV CTR. Internal and external content specialists were consulted on key areas to address.
- **Review scientific literature.** Approximately 5,000 abstracts were screened and approximately 600 relevant publications were reviewed and synthesized where appropriate. Approximately 20 previously published CDC guidelines related to HIV CTR also were summarized.
- **Obtain technical opinion.** A panel of technical specialists from public and private sectors; governmental and nongovernmental agencies; and legal, ethics, and policy fields was convened to review the recommendations.
- **Obtain user input.** Internal CDC comments, public and private provider assessments, key consultant interviews, broad external reviews, and public comments through the Federal Register were obtained.
- **Publish electronically and in hard copy.** Single copies of this report are available from CDC’s National Prevention Information Network (NPIN) website at <<http://www.cdcnpin.org>> or by calling (800) 458-5231. The guidelines are also available at the HIV Counseling, Testing, and Referral website at <<http://www.cdc.gov/hiv/ctr>>. They will be updated and posted periodically.

Similarities and Differences Between Current and Previous Guidelines

Aspects of previous CDC HIV guidelines that are unchanged include

- encouraging availability of anonymous as well as confidential HIV testing;
- ensuring that HIV testing is informed, voluntary, and consented;
- emphasizing access to testing and effective provision of test results;
- advocating routine recommendation of HIV CTR in settings (e.g., publicly funded clinics) serving clients at increased behavioral or clinical risk for HIV infection;
- recommending use of a prevention counseling approach aimed at personal risk reduction for HIV-infected persons and persons at increased risk for HIV; and
- stressing the need to provide information regarding the HIV test to all who take the test.

Differences in the new guidelines include

- giving guidance to all providers of voluntary HIV CTR in the public and private sectors;
- using an evidence-based approach to provide specific recommendations for CTR;
- underscoring the importance of early knowledge of HIV status and making HIV testing more accessible and available;
- acknowledging providers' need for flexibility in implementing the guidelines, given their particular client base, setting HIV prevalence level, and available resources;
- recommending that CTR be targeted efficiently through risk screening and other strategies; and
- addressing ways to improve the quality and provision of HIV CTR.

Advances in HIV/AIDS Prevention and Treatment Interventions

During the past 2 decades, HIV infection and severe HIV-related diseases (e.g., acquired immunodeficiency syndrome [AIDS]) have become a leading cause of illness and death in the United States. As of December 31, 2000, a total of 774,467 persons were reported with AIDS, and 448,060 of these persons had died; the number of persons living with AIDS (322,865) was the highest ever reported (35). Approximately 800,000–900,000 persons in the United States are infected with HIV, and approximately 275,000 of these persons might not know they are infected (36).

Since the last CTR guidelines were published, many advances have been made in HIV/AIDS prevention and treatment, including development of effective antiretroviral therapies that have reduced HIV-related illness and death. However, although medical treatment has improved the quality and length of life for HIV-infected persons, it cannot cure HIV disease. Furthermore, the successes of new medical therapies might have led to relaxed attitudes toward safer sex (e.g., increased incidence of unprotected anal sex by young men who have sex with men) by HIV-infected persons and uninfected persons at increased risk (36,37). Additional advances include improved understanding of HIV transmission; a wider array of HIV test technologies; effective prevention counseling approaches; and practical, beneficial referral strategies — all of which could reduce the impact of the HIV epidemic in the United States.

Early knowledge of HIV infection is now recognized as a critical component in controlling the spread of HIV infection (38). Cohort studies have demonstrated that many infected persons decrease behaviors that transmit infection to sex or needle-sharing partners once they are aware of their positive HIV status (39–46). HIV-infected persons who are unaware of their infection do not reduce risk behaviors (42,47–49). Persons tested for HIV who do not return for test results might even increase their risk for transmitting HIV to partners (50). Because medical treatment that lowers HIV viral load might also reduce risk for transmission to others (51), early referral to medical care could prevent HIV transmission in communities while reducing a person's risk for HIV-related illness and death.

The array of HIV test technologies available has expanded, possibly enhancing a person's willingness to be tested and learn his or her HIV status. HIV tests can use specimens collected by less-invasive methods (e.g., oral fluid, urine, and finger-stick

blood), in addition to serum specimens collected by venipuncture. Rapid HIV testing allows clients to receive results the same day, which is useful in urgent medical circumstances and settings where clients tend not to return for HIV test results (e.g., some STD clinics). HIV testing can also be conducted using commercially available home sample collection devices (52).

Also during the 1990s, randomized controlled trials demonstrated that, for persons at increased HIV risk, certain prevention counseling approaches can be effective in reducing high-risk behaviors and new sexually transmitted infections (5,18–21). The counseling approach used is critical to effectiveness; interactive counseling approaches directed at a client's personal risk and the situations in which risk occurs are more effective than didactic, informational approaches (5). Because personalized prevention counseling can require more provider time and training than traditional counseling approaches, providing it to everyone receiving HIV testing might not be feasible. This recognition has led to a new area of health services research — developing strategies that effectively target CTR services to persons most likely to benefit from them.

The improved health of HIV-infected persons on antiretroviral therapy, along with new test technologies and effective counseling approaches, has contributed to an improved understanding of the importance of referral to needed services. In addition, new guidelines for partner counseling and referral services (PCRS) (27) and prevention case management (28) were developed specifically for publicly funded clinics and could also be useful to providers in other settings. Specialists in the field have also identified situations in which additional counseling or psychosocial support services might benefit HIV prevention efforts. Finally, advances in several areas have led to new guidelines for preventing mother-to-child transmission (see *Revised Recommendations for HIV Screening of Pregnant Women*), treating opportunistic infections (23,53) and other sexually transmitted (29) and bloodborne diseases (30–32), and managing occupational and nonoccupational exposure and prophylaxis (54,55). These developments were considered in the formulation of the new CTR guidelines.

Despite these advances in HIV prevention and care, a substantial number of opportunities for HIV prevention through CTR are missed. At publicly funded sites, approximately 70% of persons tested received their results and information regarding the test, but fewer persons likely received HIV prevention counseling and referrals. In private settings, a lower proportion of all clients are tested, and few receive prevention counseling and referrals (56–59). In many potential testing settings (e.g., emergency departments), HIV prevention counseling and testing are not uniformly offered, and data regarding types, completion, and effectiveness of referrals are not routinely collected.

Goals of HIV CTR

- Ensure that HIV-infected persons and persons at increased risk for HIV
 - have access to HIV testing to promote early knowledge of their HIV status;
 - receive high-quality* HIV prevention counseling to reduce their risk for transmitting or acquiring HIV; and

* Delivered according to recommended protocols (for counseling, referral, and evaluation) or regulatory standards (for testing).

- have access to appropriate medical, preventive, and psychosocial support services.
- Promote early knowledge of HIV status through HIV testing and ensure that all persons either recommended or receiving HIV testing are provided information regarding transmission, prevention, and the meaning of HIV test results.

Principles of HIV CTR

Effective HIV CTR is based on the following principles:

- **Protect confidentiality of clients who are recommended or receive HIV CTR services.** Information regarding a client's use of HIV CTR services should remain private (i.e., confidential). Personal information should not be divulged to others in ways inconsistent with the client's original consent.
- **Obtain informed consent before HIV testing.** HIV testing should be voluntary and free of coercion. Informed consent before HIV testing is essential. Information regarding consent may be presented orally or in writing and should use language the client can understand. Accepting or refusing testing must not have detrimental consequences to the quality of care offered. Documentation of informed consent should be in writing, preferably with the client's signature. State or local laws and regulations governing HIV testing should be followed.

Information regarding consent may be presented separately from or combined with other consent procedures for health services (e.g., as part of a package of tests or care for certain conditions). However, if consent for HIV testing is combined with consent for other tests or procedures, the inclusion of HIV testing should be specifically discussed with the client. For a discussion of HIV testing in pregnant women, consult the guidelines for HIV screening of pregnant women (see *Revised Recommendations for HIV Screening of Pregnant Women*).

- **Provide clients the option of anonymous HIV testing.** Anonymous testing (i.e., consented voluntary testing conducted without a client's identifying information being linked to testing or medical records, including the request for testing or test results) has been used widely and effectively. Anonymous testing can benefit the health of individual persons and the public by prompting earlier entry into medical care (60). Persons who would otherwise not be tested might seek anonymous HIV testing and learn their HIV status. Consistent with public health best practices, states in which anonymous testing is not available should reconsider their policy. When the client has no clear preference regarding testing type, confidential testing (i.e., information documented in client's record) should be recommended to promote receipt of test results and linkage to follow-up counseling and referral for needed services. Clients opting for anonymous testing should be informed that the provider cannot link the client's test result to the client by name. Therefore, if the client does not return for test results, the provider will not be able to contact the client with those results.
- **Provide information regarding the HIV test to all who are recommended the test and to all who receive the test, regardless of whether prevention counseling is provided.** The information should include a description of ways in which HIV is transmitted, the importance of obtaining test results, and the meaning of HIV test results.

- **Adhere to local, state, and federal regulations and policies that govern provision of HIV services.** Laws at the local, state, and federal levels might address aspects of HIV services or regulate how services are provided to particular persons (e.g., minors). In addition, policies, local ordinances, funding source requirements, and planning processes could also affect a provider's decisions regarding which services to provide and how to provide them.
- **Provide services that are responsive to client and community needs and priorities.** Providers should work to remove barriers to accessing services and tailor services to individual and community needs. To ensure that clients find services accessible and acceptable, services can be offered in nontraditional settings (i.e., community-based or outreach settings); hours of operation can be expanded or altered; unnecessary delays can be eliminated (e.g., integrating counseling and testing for STDs/HIV with counseling and testing for hepatitis); test results can be obtained more easily (e.g., with rapid testing or by telephone in certain situations); and less-invasive specimen collection can be used (e.g., oral fluid, urine, or finger-stick blood).
- **Provide services that are appropriate to the client's culture, language, sex, sexual orientation, age, and developmental level.** These factors could affect how the client seeks, accepts, and understands HIV services. Providers should consider these factors when designing and providing HIV services to increase the likelihood of return for test results and acceptance of counseling and referral services.
- **Ensure high-quality services.** To ensure ongoing, high-quality services that serve client and community needs, providers should develop and implement written protocols for CTR and written quality assurance and evaluation procedures. Many state and local health departments have substantial expertise in providing and monitoring the quality of HIV CTR services and can be a resource to private providers or community-based or outreach settings initiating these services.

TARGETED VERSUS ROUTINELY RECOMMENDED HIV CTR

Providers in all settings (traditional and nontraditional) should ideally recommend CTR to all clients on a routine basis to ensure that all clients who could benefit from CTR receive these services. However, resources might be insufficient to permit this practice. Therefore, these guidelines contain recommendations aimed at ensuring that as many persons as possible who are HIV-infected or at risk for HIV who do not know their HIV status have access to testing, prevention counseling, and referrals.

Routinely Recommending CTR to All Clients Versus Targeting CTR to Selected Clients

Studies have documented that, in settings serving clients at increased behavioral and clinical risk for HIV infection, targeting HIV testing based on reported risk factors will miss many HIV-infected clients (61–69). However, in low prevalence settings, where most clients have minimal risk, targeting clients for HIV testing based on risk screening might be more feasible for identifying small numbers of HIV-infected persons (70). Providers should consider three factors in determining whether to recommend HIV CTR to all clients or to target only selected clients.

- Type of setting.
- HIV prevalence of the setting.
- Behavioral and clinical HIV risk of the individual clients in the setting.

Although certain types of settings serve populations at increased risk (e.g., STD clinics), others might serve individual clients at increased risk (e.g., private physicians' offices in areas of low prevalence). Individual risk can be ascertained through risk screening. Under certain circumstances — perinatal transmission, acute occupational exposure, and acute nonoccupational (i.e., high-risk sexual or needle-sharing) exposure — providers should recommend HIV CTR regardless of setting prevalence or behavioral or clinical risk, based on the respective guidelines (*Revised Recommendations for HIV Screening of Pregnant Women*,54,55).

Using Prevalence Data to Establish Service Priorities

Few data exist to define “high” and “low” HIV prevalence and describe how these definitions could help develop and prioritize HIV CTR services. A study conducted in the early 1990s for acute care hospitals with $\geq 1\%$ HIV prevalence reported that routine voluntary HIV testing of all patients within a specific age range could be a feasible way to identify a large proportion of HIV-infected patients (71). This 1% prevalence can be used as general guidance for whether to routinely recommend or target HIV counseling and testing in other settings.

The threshold of HIV prevalence that should lead to routine recommendations for HIV testing of all clients within a setting can vary within and across settings and should be set in consideration of available resources. Services could be routinely recommended in settings with HIV prevalence rates $< 1\%$ but higher than other settings in the community, according to U.S. prevalence data (72). If HIV prevalence data are outdated or unknown, providers should consult their local or state health department for assistance in determining appropriate HIV CTR strategies. Alternatively, providers could employ routine voluntary testing to obtain information on prevalence in their particular settings.

Because of the availability of antiretroviral therapy to reduce the risk for perinatal HIV transmission, all pregnant women should be recommended HIV testing regardless of setting prevalence or behavioral or clinical risk (see *Revised Recommendations for HIV Screening of Pregnant Women*).

Determining Individual HIV Risk Through Risk Screening*

A client's individual HIV risk can be determined through risk screening based on self-reported behavioral risk (Box 2) and clinical signs or symptoms. Behavioral risks include injection-drug use or unprotected intercourse with a person at increased risk for HIV. Clinical signs and symptoms include STDs, which indicate increased risk for HIV infection, or other signs or symptoms (e.g., of acute retroviral or opportunistic infections), which might suggest the presence of HIV infection. Insufficient data exist to support the efficacy of any one risk-screening approach over others (e.g., face-to-face discussion or interviews, self-administered questionnaires, computer-assisted interviews, or simple open-ended questions asked by providers) (Box 2) (61,70).

* Risk screening differs from risk assessment, which is a part of HIV prevention counseling (see HIV Prevention Counseling).

BOX 2. Examples of two risk-screening strategies to elicit client-reported HIV risks

- Open-ended question by provider, “What are you doing now or what have you done in the past that you think may put you at risk for HIV infection?”
- Screening questions* (i.e., a checklist) for use with a self-administered questionnaire, face-to-face or computer-assisted interview, or other instrument: “Since your last HIV test (if ever), have you
 - injected drugs and shared equipment (e.g., needles, syringes, cotton, water) with others?”
 - had unprotected intercourse with someone that you think might be infected (e.g., a partner who injected drugs, has been diagnosed or treated for a sexually transmitted disease [STD] or hepatitis, has had multiple or anonymous sex partners, or has exchanged sex for drugs or money)?”
 - had unprotected vaginal or anal intercourse with more than one sex partner?”
 - been diagnosed or treated for an STD, hepatitis, or tuberculosis?”
 - had a fever or illness of unknown cause?”
 - been told you have an infection related to a ‘weak immune system’?”

* Clients who respond affirmatively to ≥ 1 of these questions should be considered at increased risk for HIV.

Recommendations for Routinely Recommended and Targeted CTR by Setting and Circumstance

Decisions regarding whether to recommend routine or targeted services are based on the behavioral and clinical HIV risk of the client population in the setting, the level of HIV prevalence of the setting, and the behavioral and clinical HIV risk of individual clients (Box 3). These factors should not be used to determine recommendations for CTR in circumstances in which treatment potential exists (i.e., perinatal transmission and acute occupational or nonoccupational exposure).

Settings Serving Populations at Increased Behavioral or Clinical Risk

HIV CTR should be routinely recommended for all clients in settings where the client population is at increased behavioral or clinical risk for acquiring or transmitting HIV infection, regardless of setting prevalence (Box 4 and Figure 1). These services should be provided on-site. In these settings, clients with ongoing risk behaviors should be linked to additional HIV prevention and support services (e.g., PCRS, drug or alcohol prevention and treatment), as appropriate. HIV-infected clients should receive ongoing HIV prevention counseling applicable to their personal situation.

BOX 3. Clients who should be recommended HIV prevention counseling, testing, and referral

- All clients in settings serving client populations at increased behavioral or clinical HIV risk (regardless of setting HIV prevalence).
- Individual clients in settings with <1%* HIV prevalence who[†]
 - have clinical signs or symptoms suggesting HIV infection (e.g., fever or illness of unknown origin, opportunistic infection [including active tuberculosis disease] without known reason for immune suppression),
 - have diagnoses suggesting increased risk for HIV infection (e.g., another sexually transmitted disease [STD] or bloodborne infection),
 - self-report HIV risks (see Box 2), or
 - specifically request an HIV test.
- All clients in settings with a $\geq 1\%$ [§] HIV prevalence.[¶]
- Regardless of setting prevalence or behavioral or clinical risk,
 - all pregnant women,[¶]
 - all clients with possible acute occupational exposure, and
 - all clients with known sexual or needle-sharing exposure to an HIV-infected person.

* Or lower than other settings in the community.

[†] Constitutes risk screening; see Determining Individual HIV Risk Through Risk Screening.

[§] Or higher than other settings in the community.


[¶] Clients should be routinely recommended testing, and if risk is identified during risk screening, they should also be recommended HIV prevention counseling and referral.

BOX 4. Examples of settings that serve populations at increased behavioral or clinical risk for HIV infection

- | | |
|--|---|
| • Adolescent or school-based health clinics with high rates of sexually transmitted diseases (STD) | • Freestanding HIV test sites |
| • Clinics serving men who have sex with men | • Homeless shelters |
| • Correctional facilities, prisons, juvenile detention centers | • Outreach programs (e.g., syringe exchange programs) |
| • Drug or alcohol prevention and treatment programs | • STD clinics |
| | • Tuberculosis (TB) clinics* |

* Only persons with confirmed or suspected TB and their contacts should routinely be recommended HIV CTR.

FIGURE 1. An example of counseling, testing, and referral in settings serving populations at increased behavioral or clinical HIV risk



Low Prevalence Settings

In low prevalence settings (e.g., <1%, see Using Prevalence Data to Establish Service Priorities) where the client population is generally not at increased behavioral or clinical HIV risk, CTR should be targeted to clients based on risk screening (Figure 2). Prevention counseling and referral are recommended for persons at increased risk even if HIV testing is declined. Any client who requests HIV testing should receive it, regardless of risk. These settings likely represent most health-care settings.

High Prevalence Settings

In high prevalence settings (e.g., $\geq 1\%$), all clients should be routinely recommended HIV testing (Figure 3). Risk screening should be used to determine if HIV prevention counseling and referral should also be recommended. CTR should be provided on-site. In these settings, clients with ongoing risk behaviors identified during risk screening should be linked to additional HIV prevention and support services (e.g., PCRS and drug or alcohol prevention and treatment), as appropriate.

Circumstances For Which HIV Preventive Treatment Exists

Prophylaxis exists for a limited number of situations: perinatal transmission, acute occupational exposure, and acute nonoccupational (i.e., high-risk sexual or needle-sharing) exposure. Regardless of population risk, setting prevalence, or individual behavioral or clinical risk, voluntary HIV testing should be routinely recommended to a) all pregnant women, b) clients with acute occupational exposure, and c) clients with acute nonoccupational (e.g., high-risk sexual or needle-sharing) exposure. Regardless of whether a client receives an HIV test, HIV prevention counseling and referral should

FIGURE 2. An example of HIV counseling, testing, and referral in low prevalence settings





FIGURE 3. An example of HIV counseling, testing, and referral in high prevalence settings



target pregnant women based on risk screening and be routinely recommended to clients with either acute occupational or nonoccupational exposures. For further information, consult the respective guidelines on perinatal transmission, acute occupational exposure, and acute nonoccupational exposure (*Revised Recommendations for HIV Screening of Pregnant Women, 54,55*).

A Framework for Implementing HIV CTR

CTR are interrelated interventions that ideally should be integrated and offered in all settings. However, these guidelines acknowledge public and private providers' needs for flexibility. Certain providers might be able to offer prevention counseling but not an HIV test, whereas others might be able to offer an HIV test but not prevention counseling. Although all providers in settings serving populations at increased behavioral or clinical risk for HIV (e.g., STD clinics) should provide HIV CTR on-site, not all can. These providers should maintain clear and appropriate methods of referral to providers of prevention counseling or testing elsewhere. To ensure client referral, providers who offer HIV counseling and testing should collaborate with providers serving populations at increased risk for HIV who might not provide these services.

HIV COUNSELING

HIV counseling seeks to reduce HIV acquisition and transmission through the following:

- **Information.** Clients should receive information regarding HIV transmission and prevention and the meaning of HIV test results. Provision of information is different from informed consent.
- **HIV prevention counseling.** Clients should receive help to identify the specific behaviors putting them at risk for acquiring or transmitting HIV and commit to steps to reduce this risk. Prevention counseling can involve ≥ 1 sessions.

Information

All clients who are recommended or who request HIV testing should receive the following information, even if the test is declined:

- Information regarding the HIV test and its benefits and consequences.
- Risks for transmission and how HIV can be prevented.
- The importance of obtaining test results and explicit procedures for doing so.
- The meaning of the test results in explicit, understandable language.*
- Where to obtain further information or, if applicable, HIV prevention counseling.
- Where to obtain other services (see Typical Referral Needs).

* For example, "A negative test means no HIV was found. But if you were exposed to HIV recently — in the last 1–2 months — this test may not be able to pick that up." See Negative HIV Test Results.

In certain settings where HIV testing is offered, other useful information includes a) descriptions or demonstrations of how to use condoms correctly; b) information regarding risk-free and safer sex options (73); c) information regarding other sexually transmitted and bloodborne diseases; d) descriptions regarding the effectiveness of using clean needles, syringes, cotton, water, and other drug paraphernalia; e) information regarding drug treatment; and f) information regarding the possible effect of HIV vaccines on test results for persons participating in HIV vaccine trials (see Additional Counseling Considerations for Special Situations and Positive HIV Test Results).

For efficiency, information can be provided in a pamphlet, brochure, or video rather than a face-to-face encounter with a counselor. This approach allows the provider to focus face-to-face interactions on prevention counseling approaches proven effective with persons at increased risk for HIV infection. Information should be provided in a manner appropriate to the client's culture, language, sex, sexual orientation, age, and developmental level. Certain informational videos and large-group presentations that provide explicit information regarding correct use of condoms have proven effective in reducing new STDs (19–21,74) and could be effective in reducing HIV.

HIV Prevention Counseling

HIV prevention counseling should focus on the client's own unique circumstances and risk and should help the client set and reach an explicit behavior-change goal to reduce the chance of acquiring or transmitting HIV. HIV prevention counseling is usually, but not always, conducted in the context of HIV testing. The client-centered* HIV prevention counseling model involves two brief sessions (4,5,75), whereas other effective models are longer or involve more sessions (5–8,10,11,13–18,76–79). Regardless of the model used, in HIV prevention counseling, the counselor or provider focuses on assessing the client's personal risk or circumstances and helping the client set and reach a specific, realistic, risk-reduction goal. These guidelines avoid using the terms "pretest" and "posttest" counseling to underscore that prevention counseling is a risk-reduction process that might involve only one or >1 session.

Several models for HIV prevention counseling in conjunction with HIV testing have been developed, evaluated in controlled studies, and documented to be efficacious in changing behavior or reducing sexually transmitted infections, including individual face-to-face counseling (5,12), large- and small-group counseling with a facilitator (6,16,18,79), and video-based counseling (19). For more information regarding interventions, see *The Compendium of HIV Prevention Interventions with Evidence of Effectiveness* at <<http://www.cdc.gov/hiv/pubs/hivcompendium.pdf>>.

Client-Centered HIV Prevention Counseling

Since 1993, CDC has recommended one interactive counseling model, called client-centered HIV prevention counseling (3,4), which involves two face-to-face sessions with a provider or counselor (3–5,75,80). This model has traditionally used a two-step HIV

* Client-centered is used here to mean that the counseling sessions focus on the client's own risk circumstances, risk behaviors, and prevention needs. This term should not be confused with the more intensive, client-centered approach advocated by psychologist Carl R. Rogers, although some skills and strategies that involve the client in the prevention counseling process might be similar (Rogers CR. *Client-centered therapy: its current practice, implications, and theory*. Boston, MA: Houghton Mifflin, 1951).

testing approach in which clients are physically present at a setting for the HIV test (initial session) and then return for HIV test results (follow-up session). Each session might require 15–20 minutes (including testing and referral) for clients at increased risk for HIV, but could take only a few minutes for those at lower risk. In the first session, a personalized risk assessment* encourages clients to identify, understand, and acknowledge the behaviors and circumstances that put them at increased risk for acquiring HIV. The session explores previous attempts to reduce risk and identifies successes and challenges in these efforts. This in-depth exploration of risk allows the counselor to help the client consider ways to reduce personal risk and commit to a single, explicit step to do so. In the second session, when HIV test results are provided, the counselor discusses the test results, asks the client to describe the risk-reduction step attempted (and acknowledges positive steps made), helps the client identify and commit to additional behavioral steps, and provides appropriate referrals (e.g., to PCRS).

In one large, randomized, controlled trial, this model was reported to be

- effective at reducing high-risk sexual behaviors and new STDs (5);
- feasible to use even in busy publicly funded clinics;
- acceptable to clients, counselors, and health-care providers (80); and
- cost-effective at preventing STDs in persons at increased risk for HIV (81–83).

The model was reported to be especially effective among adolescents and persons with ongoing sexual risk behaviors (e.g., newly diagnosed STDs) (5). Although the benefits of client-centered HIV prevention counseling in reducing high-risk drug behaviors are unknown, studies have indicated that similar counseling approaches that help clients explore risks and set specific risk-reduction goals reduce risky drug use behaviors (39–41,84).

Observational studies and reviews of programs in various settings have indicated that many counselors are still unfamiliar with the specific goals of the client-centered HIV prevention counseling model (75,85) (Amy S. DeGroff, M.P.H., written communication, 2000). Because “client-centered” is sometimes misinterpreted as “face-to-face,” providers in many HIV test sites deliver face-to-face informational messages in response to a generic checklist risk assessment. This type of counseling provides advice rather than encouraging client participation or discussion of personal risk; it seldom focuses on personal goal setting. “Client-centered” can also be misinterpreted to mean that the counselor should avoid directing the session. Although attentive listening and respect for clients’ concerns are important elements of effective counseling, the primary goal of client-centered HIV prevention counseling is risk reduction. HIV prevention counseling usually requires provider training and support and ongoing quality assurance to achieve optimal benefit. Providers can contact their state health department’s HIV/AIDS program office for information on local training opportunities. For information on client-centered counseling with rapid testing, see Addressing Barriers to HIV Prevention Counseling.

* Personal risk assessment is an essential element of HIV prevention counseling in which the client and counselor work to understand and acknowledge the client’s personal risk for HIV. Risk assessment is not synonymous with risk screening (see Determining Individual Client Risk Through Risk Screening and Box 2), which helps determine which clients should be recommended HIV CTR.

Elements of HIV Prevention Counseling

Regardless of the HIV prevention counseling model used, some counseling elements have been used repeatedly in effective interventions and are recognized by many specialists as critical in counseling success (Technical Expert Panel Review of CDC HIV Counseling, Testing, and Referral Guidelines; February 18–19, 1999; Atlanta, Georgia).

The following elements should be part of all HIV prevention counseling sessions:

- **Keep the session focused on HIV risk reduction.** Each counseling session should be tailored to address the personal HIV risk of the client rather than providing a predetermined set of information. Although counselors must be willing to address problems that pose barriers to HIV risk reduction (e.g., alcohol use in certain situations), counselors should not allow the session to be distracted by the client's additional problems unrelated to HIV. Certain counseling techniques (e.g., open-ended questions [Box 5], role-play scenarios, attentive listening, and a nonjudgmental and supportive approach) can encourage the client to remain focused on personal HIV risk reduction.
- **Include an in-depth, personalized risk assessment.** Sometimes called "enhancing self-perception of risk," risk assessment allows the counselor and client to identify, acknowledge, and understand the details and context of the client's HIV risk (17,86,87). Keeping the assessment personal, instead of global, will help the client identify concrete, acceptable protective measures to reduce personal HIV risk (Box 6). The risk assessment should explore previous risk-reduction efforts and identify successes and challenges in those efforts. Factors associated with continued risk behavior that might be important to explore include using drugs or alcohol before sexual activity, underestimating personal risk, perceiving that precautionary changes are not an accepted peer norm, perceiving limited self-efficacy for successful change efforts, receiving reinforcement for frequent unsafe practices (e.g., a negative HIV test result after risk behaviors), and perceiving that vulnerability is associated with "luck" or "fate" (86–89).
- **Acknowledge and provide support for positive steps already made.** Exploring previous risk-reduction efforts is essential for understanding the strengths and challenges faced by the client in reducing risk. Support for positive steps already taken increases the clients' beliefs that they can successfully take further HIV risk-reduction steps. For some clients, simply agreeing to an HIV test is an important step in reducing risk (5,75).
- **Clarify critical rather than general misconceptions.** In most situations, counselors should focus on reducing the client's current risk and avoid discussions regarding HIV transmission modes and the meaning of HIV test results. However, when clients believe they have minimal HIV risk but describe more substantial risk, the counselor should discuss the HIV transmission risk associated with specific behaviors or activities the clients describe and then discuss lower-risk alternatives (73). For example, if clients indicate that they believe oral sex with a risky sex partner poses little or no HIV risk, the counselor can clarify that, although oral sex with an infected partner might result in lower HIV transmission risk than anal sex, oral sex is not a risk-free behavior, particularly when commonly practiced. If clients indicate that they do not need to be concerned about HIV transmission among needle-sharing partners if they use clean needles, the counselor can clarify that

BOX 5. Examples of closed-ended versus open-ended questions

Closed-ended questions, which might interfere with client-centered human immunodeficiency virus (HIV) prevention counseling	Open-ended questions, which promote client-centered HIV prevention counseling
Have you ever injected drugs? OR	What are you doing that you think may be putting you at risk for HIV infection?
Have you (for a male client) ever had sex with a man? OR	What are the riskiest things that you are doing?
Have you (for a female client) ever had sex with a bisexual man?	If your test comes back positive, how do you think you may have become infected?
Have you ever had sex when you were under the influence of alcohol or drugs?	When was the last time you put yourself at risk for HIV? What was happening then?
Do you (always) use condoms when you have sex? OR	How often do you use drugs or alcohol?
Can you always use condoms when you have sex?	How do you think drugs or alcohol influence your HIV risk?
	How often do you use condoms when you have sex?
	When/with whom do you have sex without a condom? When with a condom?
Can you always use clean works (i.e., needles, syringes, cottons, or cookers*) when you inject?	What are you currently doing to protect yourself from HIV? How is that working?
	What kinds of things do you do to protect your partner from getting infected with HIV? (for HIV-infected clients) Tell me about specific situations when you have reduced your HIV risk. What was going on that made that possible?
	How risky are your sex/needle-sharing partners? For example, have they been recently tested for HIV?

* Cottons are filters used to draw up the drug solution. Cookers include bottle caps, spoons, or other containers used to dissolve drugs.

BOX 6. Examples of global versus specific risk-reduction steps for HIV prevention counseling

Global risk-reduction steps, which are unlikely to be effective in changing behavior	Specific risk-reduction steps, which are more likely to be effective in changing behavior
Always use condoms.	Buy a condom tomorrow and try it on.
Have fewer or less risky partners.	Carry a condom next time I go out (e.g., to the bar/nightclub).
Have safer sex.	Starting today, put condoms on the night stand beside the bed.
Stop injecting drugs.	Starting tonight, require my partner to use a condom next time, or I will not have vaginal (anal) sex.
	Stop seeing (specific partner) who is seeing other people.
	Break up with (specific partner) before getting together with someone new.
	Talk honestly with (specific partner) about my HIV status and ask about his/her HIV status.
	Next time I'm out with friends and may have sex, avoid getting "high" on drugs or alcohol.
	Only kissing, etc., with (specific partner) until we both have an HIV test.
	Tomorrow, ask (specific partner) if he or she has had a recent HIV test and has been tested for other sexually transmitted diseases.
	Obtain clean works (i.e., needles, syringes, cottons, or cookers*) tomorrow so I have them before I use next time.
	Contact drug treatment center and make appointment.

* Cottons are filters used to draw up the drug solution. Cookers include bottle caps, spoons, or other containers used to dissolve drugs.

HIV can be transmitted through the cooker, cotton, or water used by several persons sharing drugs. With newly identified or uninformed HIV-infected clients, the counselor should discuss HIV transmission risks associated with specific sexual or drug-use activities, including those in which the client might not be currently engaged.

- **Negotiate a concrete, achievable behavior-change step that will reduce HIV risk.** Although the optimal goal might be to eliminate HIV risk behaviors, small behavior changes can reduce the probability of acquiring or transmitting HIV. Behavioral risk-reduction steps should be acceptable to the client and appropriate to the client's situation. For clients with several high-risk behaviors, the counselor should help clients focus on reducing the most critical risk they are willing to commit to changing. The step does not need to be a personal behavior change. For many clients, knowledge of a partner's recent HIV status (and talking with the partner about getting an HIV test) might be more critical than personal behavior changes. The step should be relevant to reducing the client's own HIV risk and should be a small, explicit, and achievable goal, not a global goal (Box 6). Identifying the barriers and supports to achieving a step, through interactive discussion, role-play modeling, recognizing positive social supports, or other methods will enhance the likelihood of success (90). Writing down the goal might be useful. For clients with ongoing risk behaviors, referral to additional prevention and support services is encouraged.
- **Seek flexibility in the prevention approach and counseling process.** Counselors should avoid a "one-size-fits-all" prevention message (e.g., "always use condoms"). Behaviors that are safe for one person might be risky for another (91). For example, unprotected vaginal intercourse might be unsafe with anonymous partners whose HIV status is unknown, but safe for uninfected persons in a mutually monogamous relationship. The length of counseling sessions will vary depending on client risk and comfort (e.g., adolescents might require more time than adults).
- **Provide skill-building opportunities.** Depending on client needs, the counselor can demonstrate or ask the client to demonstrate problem-solving strategies such as a) communicating safer sex commitments to new or continuing sex partners; b) using male latex condoms properly; c) trying alternative preventive methods (e.g., female condoms); d) cleaning drug-injection equipment if clean syringes are unavailable; or e) communicating safer drug-injection commitments to persons with whom the client shares drug paraphernalia (86,92–94).
- **Use explicit language when providing test results.** Test results should be provided at the beginning of the follow-up session. Counselors should never ask the client to guess the test results. Technical information regarding the test can be provided through a brochure or other means so the session can focus on personal HIV risk reduction for clients with negative tests and other considerations for clients with positive or indeterminate test results (see Additional Counseling Considerations for Special Situations). In-depth, technical discussions of the "window period (i.e., the time from when a person is infected until they develop detectable HIV antibody) should be avoided because they could confuse the client and diffuse the

importance of the HIV prevention message. Counselors should clarify that negative test results do not mean the client has no HIV risk and work with the client to reconsider ongoing HIV risk behaviors and the benefits of taking steps to reduce those risks. A client with ongoing risk behaviors should not be given a false sense of the safety of those behaviors (i.e., avoid statements like “whatever you were doing seems to be safe” or “continue to do whatever you are doing now”).

These counseling elements are considered necessary for high-quality counseling. Specialists in the field (Technical Expert Panel Review of CDC HIV Counseling, Testing, and Referral Guidelines; February 18–19, 1999; Atlanta, Georgia) also suggested adoption of the following:

- **Ensure that the client returns to the same counselor.** Consistency of the client and counselor relationship helps the client feel secure, reduces misunderstanding, and promotes the likelihood of effective risk reduction. Effective counseling models tended to use the same counselor for all sessions. When follow-up prevention counseling sessions must be provided by a different counselor, careful record-keeping is recommended to ensure high-quality counseling. See *The Compendium of HIV Prevention Interventions with Evidence of Effectiveness* at <<http://www.cdc.gov/hiv/pubs/hivcompendium.pdf>>.
- **Use a written protocol to help counselors conduct effective sessions.** A structured protocol outlining session goals can help keep the counselor focused on risk reduction. The protocol can include examples of open-ended questions (to help a new counselor avoid closed-ended questions) and a list of explicit risk-reduction steps (to help a new counselor avoid accepting a client’s suggestion of global risk-reduction steps) (95).
- **Ensure ongoing support by supervisors and administrators.** Supervisory support is essential for effective counseling. Training in HIV counseling approaches that focus on personal risk reduction is recommended for persons supervising counselors. Staff appraisals should acknowledge that completion of critical counseling elements has higher priority than completion of paperwork.
- **Avoid using counseling sessions for data collection.** If required, paperwork should be completed at the end of the counseling session or by staff members who are not counseling. Checklist risk assessments driven by data collection forms are detrimental to effective counseling because they can encourage even skilled counselors to use closed-ended questions, limit eye contact, and miss critical verbal and nonverbal cues. The relevance of any routinely collected data should be periodically assessed.
- **Avoid providing unnecessary information.** An emphasis on providing information might prompt counselors to miss critical HIV prevention opportunities and cause clients to lose interest. Discussion of theoretical HIV risks (e.g., sex with a person with hemophilia or needle exposures through tattoos) tends to shift the focus away from the client’s actual HIV risk situations to topics that are more “comfortable” or easy to discuss but irrelevant to the client’s risk.

Who Should Deliver Prevention Counseling

In any setting where HIV testing is provided, existing personnel can be effective counselors if they have the desire and appropriate training and employ the essential counseling elements (5,80). Advanced degrees or extensive experience are not necessary for effective HIV prevention counseling, though training is (80). Training in counseling is available (see Ensuring High-Quality HIV Prevention Counseling). In situations where primary health-care providers (e.g., physicians) might not be able to provide prevention counseling, auxiliary health professionals trained in HIV prevention counseling models can provide this service. Although peer counseling has been successful in certain situations (18), research does not support an explicit risk-reduction need or benefit to matching clients with counselors based on same or similar backgrounds, sex, ethnicity, age, or peer group for intervention efficacy (96–98). The following skills and counselor characteristics were identified by specialists in the field as important for effective HIV prevention counseling (Technical Expert Panel Review of CDC HIV Counseling, Testing, and Referral Guidelines; February 18–19, 1999; Atlanta, Georgia):

- Completion of standard training courses in client-centered HIV prevention counseling or other risk-reduction counseling models.
- Belief that counseling can make a difference.
- Genuine interest in the counseling process.
- Active listening skills.
- Ability to use open-ended rather than closed-ended questions (Box 5).
- Ability and comfort with an interactive negotiating style rather than a persuasive approach.
- Ability to engender a supportive atmosphere and build trust with the client.
- Interest in learning new counseling and skills-building techniques.
- Being informed regarding specific HIV transmission risks (73).
- Comfort in discussing specific HIV risk behaviors (i.e., explicit sex or drug behaviors).
- Ability to remain focused on risk-reduction goals.
- Support for routine, periodic, quality assurance measures.

Additional Counseling Considerations for Special Situations

- **Persons with newly identified HIV infection.** Clients with newly identified HIV infection have immediate and long-term needs. Some clients might be better prepared to receive positive test results than others. The emotional impact of hearing an HIV-positive test result might prevent clients from clearly understanding information during the session in which they receive their results. Providers should provide appropriate referrals (see Typical Referral Needs) and, when necessary, additional sessions.

When a client receives the test result, the provider should ensure that the client understands it. As part of HIV prevention counseling, providers should explicitly discuss and clarify any misconceptions regarding HIV transmission risk to partners associated with specific sexual or needle-sharing activities. Clients should be advised to refrain from donating blood, plasma, or organs. For sexually active clients who are not in mutually monogamous partnerships, providers should also address strategies to prevent other sexually transmitted or bloodborne infections (e.g., gonorrhea, syphilis, chlamydia, herpes simplex virus, human herpes virus type 8 [the virus linked to Kaposi sarcoma], hepatitis B virus, hepatitis C virus, and cytomegalovirus).

The first few months after persons learn they are HIV infected are important for accessing medical and other support services to help them obtain treatment and establish and maintain behavior changes that reduce the likelihood of transmitting the virus to others. For example, persons with ongoing risks might be referred for prevention counseling to prevent transmission to others or for prevention case management. For all newly identified clients, a follow-up appointment 3–6 months after diagnosis is recommended by some specialists (99) to assess whether clients were able to initiate medical care, minimize transmission risk to uninfected partners, and access other needed services (e.g., partner counseling and referral services). See guidance on partner counseling and referral services (27) and prevention case management (28).

- **Persons with a single, recent nonoccupational HIV exposure.** After a reported sexual, injection-drug use, or other nonoccupational exposure to HIV (55), providers should refer clients for prompt initiation of evaluation, counseling, and follow-up services. Early postexposure prophylaxis could reduce the likelihood of becoming infected with HIV, although the degree to which early treatment can prevent new infection after acute nonoccupational HIV exposure is unclear. Further guidance on nonoccupational HIV exposure is available (55).
- **Persons with indeterminate HIV test results.** Until follow-up test results are available, persons with an indeterminate result should receive information regarding the meaning of test results. HIV prevention counseling should be the same as for a person with newly identified HIV infection. Behaviors that minimize the risk for HIV transmission to sex and needle-sharing partners should be emphasized, even if the client reports no risk behaviors. Clients with repeated indeterminate test results ≥ 1 month apart are unlikely to be HIV-infected and can be provided test results in the same way as clients with negative test results, unless recent HIV exposure is suspected (see Indeterminate Test Results).
- **Persons seeking repeat HIV testing.** In addition to brief prevention counseling sessions, ongoing HIV prevention counseling aimed at personal risk reduction might be useful for persons seeking repeated HIV testing who have continued HIV risk. Counselors should encourage clients to explore alternative prevention strategies and to identify and commit to additional risk-reduction steps. Clients with ongoing risk behaviors might benefit from referral to other HIV prevention and support services because their current risk behavior might be reinforced by repeated negative HIV test results or they might view HIV testing as protective (100). More information on prevention case management is available (28) (see Ongoing Exposure).

- **Persons who use drugs.** Persons who inject drugs might also be at increased risk for acquiring HIV through unprotected sex with an HIV-infected partner (101–103). For injection-drug users (IDUs), intervention studies indicate that personalized, interactive prevention counseling models using goal-setting strategies might be effective in reducing injection-drug and sexual-risk behaviors (39–41,84). Evidence also supports the efficacy of community strategies (e.g., methadone maintenance programs or other drug treatment programs, outreach programs, and syringe exchange) to reduce new HIV infections among IDUs (104–108). Specialists in the field advocate recommending such strategies, along with individual HIV prevention counseling, to persons who inject drugs.
- **Sex or needle-sharing partners of HIV-infected persons.** Sex or needle-sharing partners of HIV-infected persons should be encouraged to have HIV prevention counseling and testing. Partners who are HIV discordant (i.e., one person is HIV-infected and the other is uninfected) should receive counseling aimed at preventing HIV transmission from the infected to the uninfected partner, including explicit discussion and clarification of any misconceptions regarding HIV transmission risk associated with specific sexual or needle-sharing activities. In addition, many HIV-discordant couples benefit from ongoing HIV prevention counseling aimed at personal risk reduction or from couples counseling that teaches safe sexual practices and proper condom use (27,109–111). Little evidence exists to conclusively support or refute whether simultaneous infection with ≥ 2 subtypes of HIV is likely to occur or, if it does, whether it is associated with more aggressive or resistant disease (112). Researchers are divided on the value of recommending consistent condom use to prevent HIV sequelae for mutually monogamous, HIV-infected partners.
- **Health-care workers after an occupational exposure.** After an occupational exposure, health-care workers should use measures to prevent transmission during the follow-up period (54). HIV-exposed health-care workers should be advised that, although HIV is infrequently transmitted through an occupational exposure, they should abstain from sex or use condoms and avoid pregnancy until they receive a negative follow-up test result. In addition, they should not donate blood, plasma, organs, tissue, or semen; if a woman is breast-feeding, she should consider discontinuing (54). Health-care workers should also be advised of the rationale for postexposure prophylaxis, the risk for occupationally acquired HIV infection from the exposure, the limitations of current knowledge of the efficacy of antiretroviral therapy when used as postexposure prophylaxis, the toxicity of the drugs involved, and the need for postexposure follow-up (including HIV testing), regardless of whether antiretroviral therapy is taken. Further guidance on occupational HIV exposure is available (54).
- **Participants in HIV vaccine trials.** HIV-vaccine-induced antibodies may be detected by current HIV tests and may cause a false-positive result. Trial participants should be advised that HIV CTR is best provided at the vaccine trial sites, the vaccine is of unknown efficacy, and HIV risk behavior can result in their becoming HIV-infected (see Positive Test Results).

Addressing Barriers to HIV Prevention Counseling

Several factors can prevent provision of high-quality HIV prevention counseling, including unavailability of trained prevention counselors at the setting in which the HIV test was conducted, client reluctance, and low rates of client return for test results. Recommended strategies for addressing these common barriers include a) providing counseling on-site, b) enhancing client acceptance of counseling by examining and improving the counseling provided, and c) considering alternate counseling methods.

Provide On-Site Counseling

Cost, lack, or turnover of trained staff members and space constraints are barriers to providing HIV prevention counseling (113). However, given the proven efficacy of prevention counseling models, in settings where HIV prevalence is high or the population served is at increased risk, the ability to provide such counseling on-site is a high priority, and efforts should be made to address and remove barriers to providing HIV prevention counseling on-site. Health educators or other auxiliary staff members trained to discuss preventive activities such as healthy eating, prenatal education, or smoking cessation could, if adequately trained, be effective HIV prevention counselors. In the interim, alternative resources should be identified, and clearly defined referrals should be made to settings that can provide high-quality prevention counseling for clients at increased HIV risk. Systems to ensure that referrals are completed should be established (see HIV Referral).

Enhance Client Acceptance of HIV Prevention Counseling

Clients who agree to HIV testing but decline HIV prevention counseling often report they lack time or already are aware of HIV transmission modes. However, experienced counselors report that clients mainly refuse counseling because they do not perceive the service to be personally beneficial (Technical Expert Panel Review of CDC HIV Counseling, Testing, and Referral Guidelines; February 18–19, 1999; Atlanta, Georgia). These counselors believe that most of these clients are concerned about a specific risk, which they would be willing to explore if the counseling seemed useful. Three of the most commonly reported barriers to the perceived usefulness of counseling are the type of counseling provided, how it is recommended, and the setting of the counseling. In settings where many clients are declining counseling, these barriers and others should be examined. The counseling might be providing information only rather than addressing personal risks. Counselors might not be offering counseling in ways appropriate to the client's culture, language, sex, sexual orientation, age, or developmental level. The setting might inhibit open discussion of risk (e.g., some outreach settings are not private). Counseling skills (e.g., attentive listening, use of open-ended questions) that allow clients to participate might have been overlooked. Even when clients at increased risk refuse counseling, use of 1–2 open-ended questions that urge clients to examine their personal situations could prompt personal exploration of risk (e.g., "What were your concerns that led you to decide to get tested today?").

Consider Alternative Methods for HIV Prevention Counseling

HIV prevention counseling models proven effective have all used face-to-face (individual or group) encounters between counselor and client and involved ≥ 2 brief sessions.

Thus, face-to-face prevention counseling is preferred for clients at increased HIV risk. Most HIV test sites use an enzyme immunoassay (EIA) and confirmatory test algorithm that requires several days for final results. The return visit for test result offers an opportunity to continue prevention counseling in a second, face-to-face meeting. However, in some settings (e.g., STD clinics, managed care organizations, and other private settings), many clients do not return for their results (50,114–116). In such settings, providers can adopt strategies that increase clients' receipt of test results, and counseling strategies might need to be adapted (117).

Telephone Counseling. Limited studies among STD clinic clients at lower risk indicated that substantially more clients learned their HIV infection status when negative test results were provided by telephone rather than in person (12,117,118). Although home sample collection provides a precedent for providing counseling by telephone to persons with either negative or positive HIV test results, the efficacy of telephone counseling in reducing HIV risk behaviors or the number of new HIV infections has not been studied. One study indicated that telephone notification of positive results was associated with delay in linkage to care (119). However, not learning positive test results at all guarantees a delay in linkage to care. Many specialists recommend that provision of HIV test results and prevention counseling by telephone be limited to clients whose results are negative (Technical Expert Panel Review of CDC HIV Counseling, Testing, and Referral Guidelines; February 18–19, 1999; Atlanta, Georgia). Also, given the known risk-reduction benefits of face-to-face counseling, lack of efficacy data on telephone counseling, and concerns regarding disinhibition (e.g., "since my test result is negative, whatever risks I am taking now may be okay"), telephone counseling should be limited to clients without known ongoing HIV risk behaviors (e.g., unprotected sex or needle-sharing with an HIV-infected [or status unknown] partner).

Single-Session Prevention Counseling with Rapid Testing. Rapid tests allow clients to receive their HIV test results the same day. This process could reduce the number of clients receiving two prevention counseling sessions. Studies of the efficacy of single HIV prevention counseling sessions for use with a rapid test are under way. Although some single-session counseling protocols have been successfully implemented in busy clinics and are well-accepted by most clients, how well a single counseling session reduces risk behaviors or the number of new HIV infections is unknown. A counseling protocol for use with a rapid test is being studied; information is available at <<http://www.cdc.gov/hiv/projects/respect-2>>. For clients with identified risk behaviors, referral or rescheduling for ongoing counseling should be considered.

Ensuring High-Quality HIV Prevention Counseling

All CTR providers should conduct routine, periodic assessments for quality assurance to ensure that the counseling being provided includes the recommended, essential counseling elements.

Supervisors must be aware of HIV prevention counseling goals and necessary counselor skills. Supervisor and administrator support of HIV counseling models that focus on personal risk reduction (distinct from provision of information) is critical to effective counseling. Quality assurance for counseling should contain the following elements:

- **Training and continuing education.** Basic training in the use of ≥ 1 of the interactive HIV prevention counseling models aimed at personal risk reduction is recommended for counselors and supervisors. Counselors should know the communities they serve and the available referral opportunities. They also might benefit from formal training on a) transmission and prevention of HIV and other sexually transmitted and bloodborne diseases, b) the natural history of HIV, c) recognition and treatment of opportunistic infections, d) new therapeutic agents used to treat HIV and AIDS, e) PCRS, f) prevention case management, and g) other HIV prevention and support services available in the community (e.g., services related to substance abuse assessment, cultural competence, adolescent concerns, domestic abuse, and health concerns for gay or lesbian clients). Additional training in specific counseling skills is also warranted (e.g., training on how to facilitate groups for counselors conducting group sessions). For training opportunities, providers or supervisors can contact their state health department's HIV/AIDS program office.
- **Supervisor observation and immediate feedback to counselors.** Direct observation of counseling sessions can help ensure that objectives are being met (80). Supervisors can perform this task periodically (with client consent). Sessions might also be audiotaped (with client consent), or counseling can be demonstrated through role-play scenarios between the counselor and supervisor. Observation and feedback should be structured, and the outcome should be constructive, not punitive. Supervisors should support positive elements of the prevention counseling session and provide specific, constructive comments regarding content areas needing improvement. Observation and feedback should be conducted regularly for routine counseling. Staff discomfort with observation typically wanes over time; many counselors report that the sessions are useful in enhancing skills. When observation is offered routinely, clients seldom refuse to participate. A suggested time frame for routine, direct observation of an HIV prevention counselor by the supervisor is twice monthly for the first 6 months, monthly for the second 6 months, and quarterly for counselors with >1 year of experience. After observation, supervisors should provide feedback to counselors quickly, preferably the same week. Observation and feedback forms used in research studies of client-centered HIV prevention counseling are available at <<http://www.cdc.gov/hiv/projects/RESPECT/default.htm>>.
- **Periodic evaluation of physical space, client flow, and time concerns.** Counseling sessions should be conducted in a private space where discussion cannot be overheard. Clients should not wait for long periods between testing and counseling, and information could be provided during waiting times (e.g., through videos). Periodic time-flow analyses or client surveys can be used to evaluate adequacy of space, client flow, and length of waiting period.
- **Periodic counselor or client satisfaction evaluations.** Evaluations of client satisfaction can ensure that counseling meets client needs. These evaluations also can provide important feedback to counselors who otherwise might not see the benefits of what they do. Evaluations can be brief. Surveys should address whether specific counseling goals were met, the type of interaction (e.g., "who talked more, the counselor or the client?"), and, when applicable, specifics of

development of the risk-reduction plan (e.g., “what was the behavior change step that you agreed to work on?”). Linking client and counselor descriptions of a particular session might be useful. Conducting such evaluations only occasionally (e.g., for 1–2 weeks once or twice a year) decreases the programmatic burden and is probably sufficient to identify problems. For more information, see Quality Assurance and Evaluation of HIV CTR Services.

- **Case conferences.** Regularly scheduled meetings of counselors allow supervisors to understand counselors’ skills and areas that need improvement and can help counselors learn techniques from their colleagues. Case conferences are an opportunity for counselors to discuss specific or problematic questions asked by clients, allowing providers to better understand the concerns facing clients who are HIV-infected or at increased risk for HIV. Case conferences can help offset counselor fatigue and “burn out” by providing a positive outlet for dealing with difficult situations. Discussion might focus on a hard-to-address client or specific elements (e.g., developing acceptable and practical risk-reduction plans with clients who deny the magnitude of their HIV risk). Frequency of case conferences should be balanced with client volume, with efforts made to meet at least monthly.

HIV TESTING

Characteristics and Applications of HIV Test Technologies

Only FDA-approved HIV tests should be used for diagnostic purposes. Routine screening in the United States for HIV-2 and HIV-1 group O infections is not generally recommended unless geographic, behavioral, or clinical information indicates that infection with these strains might be present. Several HIV test technologies have been approved by FDA for diagnostic use in the United States. These tests enable testing of different fluids (i.e., whole blood, serum, plasma, oral fluid, and urine) (Table). The available technologies

- enable specimen collection procedures that are less invasive and more acceptable than venipuncture, thus helping expand HIV testing into nontraditional settings (with home sample collection tests, oral fluid tests, and urine-based tests) (25);
- enable provision of HIV test results during a single visit at the time of testing (with rapid tests) (120); and
- increase the convenience of HIV testing (with home sample collection tests) (52).

The decision to adopt a particular test technology in a clinical or nontraditional setting should be based on several factors, including

- accuracy of the test,
- client preferences and acceptability,
- likelihood of client returning for results,
- cost and mechanism for provider reimbursement,
- ease of sample collection,

TABLE. Performance attributes and potential applications of HIV test technologies approved by the U.S. Food and Drug Administration (FDA) for diagnostic use

Test type	Specimen (mode of collection)	Test complexity*	Screening; confirmatory	Strains detected [†]	Provision of results	Advantages	Potential settings
Standard HIV test	Serum or plasma (phlebotomy)	High	Enzyme immunoassay (EIA); Western blot or immunofluorescence assay (IFA)	HIV-1 and HIV-2	HIV negative: Test result at return visit (typically a few days to 1–2 weeks) HIV positive: Confirmed result at return visit	<ul style="list-style-type: none"> • High sensitivity • Rare false-positives • High-volume processing • Utility for testing for other conditions (e.g., sexually transmitted diseases [STDs]) 	<ul style="list-style-type: none"> • Blood screening • Various settings and populations
Rapid test	Serum, plasma, whole blood (phlebotomy, finger stick)	Moderate [§]	Rapid EIA; Western blot/IFA [¶]	HIV-1	HIV negative: Test result at time of testing (typically 10–60 minutes) HIV positive: Preliminary positive test result at time of testing; ** confirmed result at return visit	<ul style="list-style-type: none"> • Convenience • Increased receipt of test results • Use in urgent medical circumstances (e.g., postexposure prophylaxis) 	<ul style="list-style-type: none"> • Settings with low return rates • Perinatal/labor and delivery for prophylaxis • Health-care settings for decisions regarding postexposure prophylaxis
Home sample collection test ^{††}	Dried blood spot (finger stick)	High	EIA; Western blot/IFA	HIV-1	HIV negative: Test result when client telephones (typically 3–7 days) HIV positive: Confirmed result when client telephones	<ul style="list-style-type: none"> • Convenience • Anonymity • Privacy • Conservation of public resources 	<ul style="list-style-type: none"> • Outreach settings • Community-based settings • Syringe exchange programs • Rural areas • Settings serving clients not at increased risk • Home

TABLE. (Continued) Performance attributes and potential applications of HIV test technologies approved by the U.S. Food and Drug Administration (FDA) for diagnostic use

Test type	Specimen (mode of collection)	Test complexity*	Screening; confirmatory	Strains detected†	Provision of results	Advantages	Potential settings
Oral fluid test	Oral mucosal transudate (oral fluid collection device)	High	EIA; Oral mucosal transudate Western blot	HIV-1	HIV negative: Test result at return visit (typically 1–2 weeks) HIV positive: Confirmed result at return visit	<ul style="list-style-type: none"> • Noninvasive • Nontechnical collection • No venipuncture • Decreased infectious hazard • Utility in nonclinical settings 	<ul style="list-style-type: none"> • Outreach settings • Community-based settings • Syringe exchange programs • Drug treatment centers • Adolescent and school-based clinics and university health centers
Urine-based test	Urine (Urine cup)	High	EIA; Urine Western blot	HIV-1	HIV negative: Test result at return visit (typically 1–2 weeks) HIV positive: Test result at return visit; further confirmation by blood sample recommended because of lower specificity of urine Western blot compared with serum-based Western blot/IFA	<ul style="list-style-type: none"> • Noninvasive • Nontechnical collection • No venipuncture • Decreased infectious hazard • Utility in nonclinical settings • Utility for testing for other conditions (e.g., STDs) 	<ul style="list-style-type: none"> • Outreach settings • Community-based settings • Syringe exchange programs • Drug treatment centers • Adolescent and school-based clinics and university health centers

* Complexity of specimen collection and testing as categorized by the Clinical Laboratory Improvement Amendments (CLIA). (Schochetman G, George JR, eds. AIDS testing: a comprehensive guide to technical, medical, social, legal, and management issues. 2 ed. New York, NY: Springer-Verlag, 1994.)

† All licensed enzyme immunoassays (EIAs) detect HIV-1, but not all detect HIV-2. EIAs that can detect HIV-1 and HIV-2 are required for blood donor screening and are recommended for diagnostic screening only where HIV-2 infection is likely. No licensed confirmatory test exists for HIV-2. Although current tests detect most HIV-1 group O infections, few detect all such infections.

‡ The one rapid test licensed by FDA, Abbott Murex Single Use Diagnostic System (SUDS) HIV-1 test (Abbott Laboratories, Inc., Abbott Park, Illinois) is classified as a moderate-complexity test and requires on-site laboratory testing capability. Future rapid tests could be classified by CLIA as “waived” and not require on-site laboratory testing capability, depending on the expertise required to perform the test correctly.

¶ Future rapid tests might be able to be confirmed with a second rapid test to provide an immediate test result with high sensitivity, specificity, and predictive value comparable with EIA/Western blot (Stetler HC, Granade TC, Nunez CA, et al. Field evaluation of rapid HIV serologic tests for screening and confirming HIV-1 infection in Honduras. AIDS 1997;11:369–75).

** Information on providing “preliminary” positive test results from a single rapid test is available elsewhere (CDC. Update: HIV counseling and testing using rapid tests—United States, 1995. MMWR 1998;47:211–5).

†† Home sample collection is different from home-use testing. FDA has approved home sample collection, but not home-use HIV test kits (Kassler WJ. Advances in HIV testing technology and their potential impact on prevention. AIDS Educ Prev 1997;9[suppl B]:27–40).

- complexity of laboratory services required for the test,
- availability of trained personnel, and
- FDA approval of the test.

Home Testing Versus Home Sample Collection

FDA has not approved home-use HIV test kits, which allow consumers to purchase a test kit, collect a sample in private, and interpret their own HIV test results in a few minutes. The Federal Trade Commission has warned that some home-use HIV test kits, many of which are available on the Internet and in the "gray" market (i.e., unauthorized imports), supply inaccurate results (121). These tests are different from FDA-approved home sample collection kits (52), which allow consumers to purchase test kits, collect a sample in private, send the sample to a laboratory for testing, and telephone for their HIV test result, counseling, and referral.

HIV-2 and HIV-1 Group O Infections

Although most HIV infections in the United States are of HIV-1 group B subtype, current EIAs can accurately identify infections with nearly all non-B subtypes and many infections with group O HIV subtypes (122). Infections with HIV-2 and HIV-1 group O are rare in the United States (123,124), and routine screening for these subtypes is not generally recommended as part of diagnostic testing except in areas where several such infections have been identified. Routine screening for HIV-2 might be appropriate in certain populations where potential risk for HIV-2 infection is higher (e.g., in areas where West African immigrants have settled) (125). Since June 1992, FDA has recommended routine screening for antibody to HIV-2 (in addition to HIV-1) for all blood and plasma donations (125). Clients with clinical, epidemiologic, or laboratory history that suggests HIV infection and negative or indeterminate HIV-1 screening tests should receive further diagnostic testing to rule out HIV infection, potentially including testing for HIV-1 non-B subtypes (122) and HIV-2 (125).

Other Test Uses

Viral load and HIV-1 p24 antigen assays are not intended for routine diagnosis but could be used in clinical management of HIV-infected persons in conjunction with clinical signs and symptoms and other laboratory markers of disease progression. Although HIV-1 p24 antigen assays are used for routine screening in blood and plasma centers, routine use for diagnosing HIV infection has been discouraged because the estimated average time from detection of p24 antigen to detection of HIV antibody by standard EIA is 6 days, and not all recently infected persons have detectable levels of p24 antigen (126).

Interpreting HIV Test Results

Standard Testing Algorithm

HIV-1 testing consists of initial screening with an EIA to detect antibodies to HIV-1. Specimens with a nonreactive result from the initial EIA are considered HIV-negative unless new exposure to an infected partner or partner of unknown HIV status has

occurred. Specimens with a reactive EIA result are retested in duplicate. If the result of either duplicate test is reactive, the specimen is reported as repeatedly reactive and undergoes confirmatory testing with a more specific supplemental test (e.g., Western blot or, less commonly, an immunofluorescence assay [IFA]). Only specimens that are repeatedly reactive by EIA and positive by IFA or reactive by Western blot are considered HIV-positive and indicative of HIV infection. Specimens that are repeatedly EIA-reactive occasionally provide an indeterminate Western blot result, which might represent either an incomplete antibody response to HIV in specimens from infected persons or nonspecific reactions in specimens from uninfected persons (127). Although IFA can be used to resolve an indeterminate Western blot sample, this assay is not widely used. Generally, a second specimen should be collected ≥ 1 month later and retested for persons with indeterminate Western blot results. Although much less commonly available, nucleic acid testing (e.g., viral RNA or proviral DNA amplification method) could also help resolve an initial indeterminate Western blot in certain situations. A small number of tested specimens might provide inconclusive results because of insufficient quantity of specimen for the screening or confirmatory tests. In these situations, a second specimen should be collected and tested for HIV infection.

Modified Testing Algorithms

FDA has licensed only one rapid test, but modified testing algorithms are anticipated when additional rapid HIV tests are approved. If ≥ 2 sensitive and specific rapid HIV tests became available, one positive rapid test could be confirmed with a different rapid test. This combination has provided positive predictive value compared with the EIA/Western blot or IFA algorithm (128). However, no such algorithms have been adequately assessed or approved for diagnostic use in the United States.

Positive HIV Test Results

An HIV test should be considered positive only after screening and confirmatory tests are reactive. A confirmed positive test result indicates that a person has been infected with HIV. False-positive results when both screening and confirmatory tests are reactive are rare. However, the possibility of a mislabeled sample or laboratory error must be considered, especially for a client with no identifiable risk for HIV infection. HIV-vaccine-induced antibodies may be detected by current tests and may cause a false-positive result. Persons whose test results are HIV-positive and who are identified as vaccine trial participants should be encouraged to contact or return to their trial site or an associated trial site for HIV CTR services.

Negative HIV Test Results

Because a negative test result likely indicates absence of HIV infection (i.e., high negative predictive value), a negative test need not be repeated in clients with no new exposure in settings with low HIV prevalence. For clients with a recent history of known or possible exposure to HIV who are tested before they could develop detectable antibodies (129,130), the possibility of HIV infection cannot be excluded without follow-up testing (29). A false negative result also should be considered in persons with a negative HIV-1 test who have clinical symptoms suggesting HIV-1 infection or AIDS. Additional testing for HIV-2 and HIV-1 group O infection might be appropriate for these persons.

Indeterminate HIV Test Results

Most persons with an initial indeterminate Western blot result who are infected with HIV-1 will develop detectable HIV antibody within 1 month (127, 131, 132). Thus, clients with an initial indeterminate Western blot result should be retested for HIV-1 infection ≥ 1 month later.* Persons with continued indeterminate Western blot results after 1 month are unlikely to be HIV-infected and should be counseled as though they are not infected unless recent HIV exposure is suspected.

Nucleic acid tests for HIV DNA or RNA exist, but are not approved by FDA for diagnostic purposes and are not generally recommended for resolving indeterminate Western blot results. However, in consultation with clinical and laboratory specialists, nucleic acid testing (if available) might also be useful for determining infection status among persons with an initial indeterminate Western blot result.

Informing Clients of Test Results

Because low rates of return for test results occur in many settings offering HIV CTR (133), providers should work to ensure that clients tested for HIV infection receive their test results, particularly HIV-infected clients who might benefit from earlier entry into care and initiation of antiretroviral therapy. Reducing barriers to testing can maximize the number and proportion of persons tested for HIV who receive their test results in a timely manner (see Addressing Barriers to HIV Testing). Adoption of new HIV test technologies and alternative methods of providing HIV-negative test results should be considered when face-to-face rates of return for test results are low. Strict confidentiality of the receipt of the HIV test and the HIV test result must be maintained, regardless of the method used. Providers unable to locate clients who do not return for test results should seek support from their local or state health department.

Because knowledge of HIV status is a critical HIV prevention strategy and essential for entry into care, providers should stress to clients the importance of returning to receive their test results and establish a plan for doing so with the client. Reminder systems might be useful. Using alternate HIV test technologies might increase the percentage of tested persons who learn their HIV status.

Providing Test Results by Telephone

Many clinicians routinely notify clients of negative test results for various diseases and conditions by means other than face-to-face (e.g., by telephone). They also ask clients to return to discuss positive test results that might indicate potential life-threatening illnesses. This strategy can also be applied, under limited circumstances, to notifying clients of their HIV test results. Face-to-face provision of HIV test results is strongly encouraged for HIV-infected clients and HIV-uninfected clients at increased risk who might benefit from HIV prevention counseling and referral to medical, preventive, and support services. Providing uninfected clients who are not at increased risk the option of receiving HIV test results and counseling by telephone — with the understanding that provider assurance of client confidentiality is of paramount importance — might be appropriate. Limited research indicates that offering clients the option of contacting

* Studies on repeat testing for persons with indeterminate Western blot results have not included pregnant women (see *Revised Recommendations for HIV Screening of Pregnant Women*).

the provider by telephone to receive negative HIV results might increase rates of receipt of results, satisfy client preferences for options, and preserve setting resources without apparent adverse consequences (52,117,118). Although no published research exists regarding use of telephones for providing positive HIV test results with most HIV test technologies, limited experience exists on using this method to provide HIV-positive test results for home sample collection testing (52).

Providing Test Results During the Initial Visit Through Rapid Tests

More clients receive their HIV test results with rapid tests because results can be provided at the testing visit (120). Rapid test technology could be useful in urgent medical circumstances (e.g., when decisions must be made regarding postexposure prophylaxis) and in nontraditional settings with low return rates (e.g., community-based or outreach settings).

During the initial visit, the provider can definitively tell clients who have had a single rapid HIV test with negative results that they are not infected (120), except when retesting might be indicated because of recent known or possible exposure to HIV. A reactive rapid HIV test result should be considered preliminary until the completion of confirmatory testing, and results should be carefully communicated to the client because of the possibility of a false-positive result.

The likelihood that a positive screening test truly indicates the presence of HIV infection decreases as HIV prevalence in the tested population becomes lower. Therefore, false-positive HIV test results are more likely in settings where the tested population prevalence is lower than in settings where the tested population prevalence is higher. When a preliminary, positive rapid test is explained to clients, phrases like “a good chance of being infected” or “very likely infected” can be used to indicate the likelihood of HIV infection and qualified based on the HIV prevalence in the setting and the client’s individual risk (120). Further testing is always required to confirm a reactive screening test result.

Follow-up Testing in Clients with Negative HIV Test Results

A negative HIV test usually indicates the absence of HIV infection (29). Because recent infection cannot be excluded without follow-up testing (see Negative HIV Test Results), the appropriate timing and frequency for follow-up testing among clients with negative HIV test results has not been firmly established. Providers should consider the following factors related to individual client needs when recommending the timing and frequency for follow-up HIV testing:

- Timing of the last potential exposure.
- Probability of HIV infection given type of exposure.
- Presence or likelihood of ongoing risk behavior.
- Likelihood of returning for follow-up HIV testing, prevention counseling, and referral.
- Client anxiety.
- Provider and client relationship.
- Resource constraints.

Recent Exposure

Follow-up testing might be appropriate for clients who have negative test results but who have not had time to develop detectable antibody after a recent documented occupational (54) or nonoccupational (sexual or needle-sharing) (55) exposure to HIV-infected persons or persons at increased risk for HIV with unknown HIV status. The timing of follow-up testing should provide assurance that the exposure did not lead to infection. Follow-up testing should be conducted in a timely manner so clients identified as HIV-infected can receive appropriate antiretroviral treatment and prevention and support services as soon as possible.

Single Possible or Known Exposure

Most infected persons will develop detectable HIV antibody within 3 months of exposure (126). If the initial negative HIV test was conducted within the first 3 months after exposure, repeat testing should be considered ≥ 3 months after the exposure occurred to account for the possibility of a false-negative result. If the follow-up test is nonreactive, the client is likely not HIV-infected. However, if the client was exposed to a known HIV-infected person or if provider or client concern remains, a second repeat test might be considered ≥ 6 months from the exposure. Rare cases of seroconversion 6–12 months after known exposure have been reported (134). Extended follow-up testing beyond 6 months after exposure to account for possible delayed seroconversion is not generally recommended and should be based on clinical judgment and individual clients needs (54).

Ongoing Exposure

Persons with continued HIV risk behavior pose a special challenge for follow-up testing. In some settings, clients with ongoing risk represent a substantial proportion of those receiving HIV CTR. In most circumstances, follow-up HIV testing should be recommended periodically for clients with ongoing risk behavior. Follow-up testing would monitor the client's HIV status, but also promote continued client contact, opportunities for HIV prevention counseling (see Additional Counseling Considerations for Special Situations), and referral to additional preventive and support services.

No Identifiable Risk

In general, persons with no recent identifiable risk for HIV infection should receive additional HIV prevention counseling and follow-up testing when requested. Efforts should be made to understand why these clients repeatedly seek follow-up testing. These clients should be considered for in-depth prevention counseling and referral to support services, where appropriate.

Special Considerations

General recommendations for follow-up testing might not be applicable in all circumstances. In certain circumstances (e.g., when persons are simultaneously exposed to hepatitis C virus and HIV [54] and when persons have received HIV vaccines), guidance should be provided only after consultation with specialists.

Addressing Barriers to HIV Testing

Knowledge of HIV infection status can benefit the health of individual persons and the community. Thus, HIV testing should be as convenient as possible to promote client knowledge of HIV infection status. Efforts should be made to remove or lower barriers to HIV testing by ensuring that

- testing is accessible, available, and responsive to client and community needs and priorities;
- anonymous and confidential HIV testing are available;
- the testing process considers the client's culture, language, sex, sexual orientation, age, and developmental level; and
- confidentiality is maintained (see Principles of HIV Counseling, Testing, and Referral).

Acceptance of HIV testing is reportedly lower when clients have been tested previously and are fearful of their ability to cope with their test results (112,113). Testing is more likely to be accepted when

- clients perceive their own HIV risk and acknowledge behaviors placing them at increased risk (135);
- testing is voluntary and routinely offered to clients rather than clients having to request it (113,136);
- protections for client confidentiality are in place (113,137);
- anonymous testing is available (113,138);
- alternate HIV test technologies are offered to clients (26);
- providers recommend testing as part of appropriate medical care (139,140); and
- providers (141) and clients (113) perceive HIV counseling and testing to be beneficial for early diagnosis and prevention purposes.

Ensuring High-Quality Testing

Testing activities should be coordinated with state and local laboratories to ensure high-quality HIV testing through proper specimen collection, storage, and transport. Laboratory errors most often occur in the preanalytic (i.e., specimen collection, labeling, transporting, processing, and storing) and postanalytic steps of testing (i.e., results validation and reporting) (142–144) rather than during the test itself. Laboratories performing HIV testing must be enrolled in proficiency testing programs and conduct activities in accordance with regulatory standards outlined by the Clinical Laboratory Improvement Amendments (CLIA) of 1988 (145). Many states have additional licensing requirements for laboratories conducting diagnostic HIV testing.

HIV REFERRAL

Definition of Referral

In the context of HIV prevention counseling and testing, referral is the process by which immediate client needs for care and supportive services are assessed and prioritized and clients are provided with assistance (e.g., setting up appointments, providing transportation) in accessing services. Referral should also include follow-up efforts necessary to facilitate initial contact with care and support service providers.

In this context, referral does not include ongoing support or management of the referral or case management. Case management is generally characterized by an ongoing relationship with a client that includes comprehensive assessment of medical and psychosocial support needs, development of a formal plan to address needs, substantial assistance in accessing referral services, and monitoring of service delivery.

Typical Referral Needs

Clients should be referred to services that are responsive to their priority needs and appropriate to their culture, language, sex, sexual orientation, age, and developmental level. Examples of these services include

- **Prevention case management.** Clients with multiple and complex needs that affect their ability to adopt and sustain behaviors to reduce their risk for transmitting or acquiring HIV should receive or be referred for prevention case management services, including ongoing prevention counseling (28). Prevention case management can help coordinate diverse referral and follow-up concerns.
- **Medical evaluation, care, and treatment.** HIV-infected clients should receive or be referred to medical services that address their HIV infection (including evaluation of immune system function and screening, treatment, and prevention of opportunistic infections) (23,29–32,53). Screening and prophylaxis for opportunistic infections and related HIV-conditions (e.g., cervical cancer) are important for HIV-infected persons. In addition, coinfection with HIV and communicable diseases (e.g., TB, STDs, and hepatitis) can, if untreated, pose a risk to susceptible community members. Thus, providers of HIV testing should be familiar with appropriate screening tests (e.g., TB), vaccines (e.g., hepatitis A and B), STD and prophylactic TB treatment, and clinical evaluation for active TB disease to ensure that these communicable diseases are identified early and appropriate clinical referrals are made, even if clients forego outpatient HIV treatment.
- **Partner counseling and referral services.** Persons with HIV-positive test results should receive or be referred to services to help them notify their sex or injection-drug-equipment-sharing partners or spouses regarding their exposure to HIV and how to access CTR. Guidelines for PCRS are available (27).
- **Reproductive health services.** Female clients who are pregnant or of childbearing age should receive or be referred to reproductive health services. HIV-infected pregnant women should be referred to providers who can provide prevention counseling and education, initiate medical therapy to prevent perinatal transmission, and provide appropriate care based on established treatment

guidelines (see *Revised Recommendations for HIV Screening of Pregnant Women*).

- **Drug or alcohol prevention and treatment.** Clients who abuse drugs or alcohol should receive or be referred to substance or alcohol abuse prevention and treatment services.
- **Mental health services.** Clients who have mental illness, developmental disability, or difficulty coping with HIV diagnosis or HIV-related conditions should receive or be referred to appropriate mental health services.
- **Legal services.** Clients who test positive should be referred to legal services as soon as possible after learning their test result for counseling on how to prevent discrimination in employment, housing, and public accommodation by only disclosing their status to those who have a legal need to know.
- **STD screening and care.** Clients who are HIV-infected or at increased risk for HIV are at risk for other STDs and should receive or be referred for STD screening and treatment (146).
- **Screening and treatment for viral hepatitis.** Many clients who are HIV-infected or at increased risk for HIV are at increased risk for acquiring viral hepatitis (A, B, and C). Men who have sex with men and IDUs should be vaccinated for hepatitis A. All clients without a history of hepatitis B infection or vaccination should be tested for hepatitis B, and if not infected, should receive or be referred for hepatitis B vaccination. In addition, clients who inject drugs should be routinely recommended testing for hepatitis C. All clients who are infected with hepatitis viruses should be referred for appropriate treatment. Further guidance is available (30,32).
- **Other services.** Clients might have multiple needs that can be addressed through other HIV prevention and support services (e.g., assistance with housing, food, employment, transportation, child care, domestic violence, and legal services). Addressing these needs can help clients access and accept medical services and adopt and maintain behaviors to reduce risk for HIV transmission and acquisition. Clients should receive referrals as appropriate for identified needs.

Implement and Manage Referral Services

Clients should receive help accessing and completing referrals, and completion of referrals should be verified. In the context of HIV prevention counseling and testing, the following elements should be considered essential to the development and delivery of referral services (99).

Assess Client Referral Needs

Providers should consult with the client to identify essential factors that a) are likely to influence the client's ability to adopt or sustain behaviors to reduce risk for HIV transmission or acquisition or b) promote health and prevent disease progression. Assessment should include examination of the client's willingness and ability to accept and complete a referral. Service referrals that match the client's self-identified priority needs are more likely to be successfully completed than those that do not (147). Priority should be placed

on ensuring that HIV-infected clients are assessed for referral needs related to medical care, PCRS, and prevention and support services aimed at reducing the risk for further transmission of HIV. When a provider cannot make appropriate referrals or when client needs are complex, clients should be referred to a case management system.

Plan the Referral

Referral services should be responsive to clients' needs and priorities and appropriate to their culture, language, sex, sexual orientation, age, and developmental level. In consultation with clients, providers should assess and address any factors that make completing the referral difficult (e.g., lack of transportation or child care, work schedule, cost). Research has indicated that referrals are more likely to be completed if services are easily accessible to clients (147).

Help Clients Access Referral Services

Clients should receive information necessary to successfully access the referral service (e.g., contact name, eligibility requirements, location, hours of operation, telephone number). Research has indicated that providing assistance (e.g., setting an appointment, addressing transportation needs) for some clients promotes completion of referrals (148). Clients must give consent before identifying information to help complete the referral can be shared. Outreach workers and peer counselors/educators can be an important and effective resource to help clients identify needs and plan successful referrals (149). Referrals are more likely to be completed after multiple contacts with outreach workers (147).

Document Referral and Follow-Up

Providers should assess and document whether the client accessed the referral services. If the client did not, the provider should determine why; if the client did, the provider should determine the client's degree of satisfaction. If the services were unsatisfactory, the provider should offer additional or different referrals. Documentation of referrals made, the status of those referrals, and client satisfaction with referrals should help providers better meet the needs of clients. Information obtained through follow-up of referrals can identify barriers to completing the referral, responsiveness of referral services in addressing client needs, and gaps in the referral system.

Ensure High-Quality Referral Services

Providers of referral services should know and understand the service needs of their clients, be aware of available community resources, and be able to provide services in a manner appropriate to the clients' culture, language, sex, sexual orientation, age, and developmental level, given local service system limitations.

Education and Support of Staff Members

Staff members providing referral services must understand client needs, have skills and resources to address these needs, have authority to help the client procure services, and be able to advocate for clients.

Training and Education. Providers should ensure that staff members receive adequate training and continuing education to implement and manage referrals. Training

and education should address resources available and methods for managing referrals, as well as promote understanding of factors likely to influence the client's ability and willingness to use a referral service (e.g., readiness to accept the service, competing priorities, financial resources). Referrals are more likely to be completed when a provider is able to correctly evaluate a client's readiness to adopt risk-reducing behaviors (147). Research has indicated that cross-training increases knowledge and understanding of community resources among providers and can indicate gaps in services (148).

Authority. Staff members providing referrals must have the authority necessary to accomplish a referral. Supervisors must ensure that staff members understand referral policy and protocol and have the necessary support to provide referrals. This requires the authority of one provider to refer to another (e.g., through memoranda of agreement) or to obtain client consent for release of medical or other personal information.

Advocacy. Staff members who negotiate referrals must possess knowledge and skills to advocate for clients. Such advocacy can help clients obtain services by mediating barriers to access to services and promoting an environment in which providers are better informed regarding the needs and priorities of their clients.

Provider Coordination and Collaboration

Providers should develop and maintain strong working relationships with other providers and agencies that might be able to provide needed services. Providers who offer HIV prevention counseling and testing but not a full range of medical and psychosocial support services should develop direct, clearly delineated arrangements with other providers who can offer needed services. Coordination and collaboration promotes a shared understanding of the specific medical and psychosocial needs of clients requiring services, current resources available to address these needs, and gaps in resources.

Memoranda of agreement or other forms of formal agreement are useful in outlining provider/agency relationships and delineating roles and responsibilities of collaborating providers in managing referrals. When confidential client information is shared between coordinating providers, such formal agreements are essential. These agreements should be reviewed periodically and modified as appropriate.

Referral Resources

Knowledge of available support services is essential for successful referrals. When referral resources are not available locally, providers should identify appropriate resources and link clients with them. A resource guide should be developed and maintained to help staff members make appropriate referrals (Box 7). Information regarding community resources can be obtained from local health planning councils, consortia, and community planning groups. Local, state, and national HIV/AIDS information hotlines or websites (e.g., NPIN), community-based health and human service providers, and state and local public health departments can also provide information.

HIV CTR SERVICES IN NONTRADITIONAL SETTINGS

CTR should be provided in community-based and outreach settings as well as clinical settings. Data from publicly supported CTR programs have indicated that doing so could promote use of these services by persons at increased risk for HIV. When HIV CTR are not readily available, accessible, or acceptable, persons at increased risk might not take

BOX 7. Contents of a referral resource guide

For each resource, the referral resource guide should specify the following:

- Name of the provider or agency
- Range of services provided
- Target population
- Service area(s)
- Contact names and telephone and fax numbers, street addresses, e-mail addresses
- Hours of operation
- Location
- Competence in providing services appropriate to the client's culture, language, sex, sexual orientation, age, and developmental level
- Cost for services and acceptable methods of payment
- Eligibility
- Application materials
- Admission policies and procedures
- Directions, transportation information, and accessibility to public transportation
- Client satisfaction with services

advantage of them. Expanding CTR into nontraditional settings can be accomplished through partnership with community-based service providers and use of new, FDA-approved HIV test technologies that offer portability, less-invasive sample collection, less-complex sample collection and processing, and reduced biohazard. To ensure effective CTR that is responsive to client needs, providers should develop and implement written quality assurance protocols and procedures specifically for services provided in nontraditional settings.

Privacy and Confidentiality

Ensuring clients' privacy and confidentiality during CTR is essential, but could present unique challenges in some nontraditional settings. Confidentiality can more easily be breached in settings where clients and providers can be seen or heard by others. Suggested strategies for maintaining privacy and confidentiality in nontraditional settings include the following:

- Use a separated area in a mobile van.
- Use rooms with locking doors.

- Mark a specific room with a “do not disturb” or “occupied” sign.
- Designate an area in the setting that provides physical privacy.
- In parks and similar locations, seek areas with as much privacy as possible.
- Provide counseling and testing services in the client’s home or other secure setting.
- Have clients return to the setting to receive test results and counseling and referral.

Informed Consent

Staff members providing CTR services should be sensitive to barriers that can interfere with obtaining true informed consent, including alcohol and drug use, mental illness, and peer pressure in venues where persons congregate or socialize. Suggested strategies for obtaining informed consent in nontraditional settings include the following:

- Schedule an appointment to test at a later date/time.
- Follow up at a later time with the client if contact information is available.
- Read the informed consent form to the client.
- Use verbal prompts to ensure that the client understands information in the informed consent form.

Counseling

Staff members working in community-based and other nontraditional settings should know and use risk-screening strategies to determine whether HIV prevention counseling should be recommended. Staff members should be trained in HIV prevention counseling or other approaches aimed at personal HIV risk reduction. When appropriate (e.g., among IDUs), information regarding other STDs and bloodborne diseases should be incorporated into the counseling sessions (29,30).

Testing

The decision to offer HIV testing in nontraditional settings should be based on several factors, including availability of resources and feasibility of providing test results and follow-up. In some cases, referral to other providers is appropriate. The selection of a specific HIV test technology should be based on logistical issues (e.g., field conditions related to collection, transport, and storage of specimens; worker safety; and the likelihood that clients will receive HIV test results). Providers must understand the extent to which field conditions can affect specimens (e.g., extreme temperatures or time lapse from collection to processing). Test specimens should be collected, stored, and transported according to manufacturer instructions.

Provision of Test Results

Clear protocols for provision of test results and prevention counseling should be developed. The following strategies might be useful in ensuring the provision of results in nontraditional settings:

- Provide a telephone number that clients can call to receive test results.
- Make an appointment with the client at the time of testing to receive results.
- Provide incentives (e.g., food certificates, hygiene kits, food).
- Return to a site on a regularly scheduled basis.
- Provide reminders when contact information is available.

Referral

Staff members working in community-based and outreach settings should be trained to implement and manage referrals. Providers should establish appropriate collaborative relationships for referrals. Arranging for PCRS staff members or case managers to be available to clients at the time test results are provided might help promote referral.

Record Keeping

Maintaining the confidentiality of client records is critical. Providers should develop written protocols for record keeping that address transport of client records to and from outreach venues. Strategies to maintain confidentiality of client records in nontraditional settings include the following:

- Return all client records to the office immediately after the CTR session.
- Use codes or unique identifiers rather than client names.
- Store all records in a secured area (e.g., locked file drawers).
- Provide option of anonymous counseling and testing as well as confidential counseling and testing.
- Verify identity of client (e.g., match client signature with that provided for informed consent or check identification card) when providing test results.
- Store paperwork in a lockbox while in outreach settings.
- Password protect and encrypt electronically stored client records.

Where allowed by state/local statute, clients can choose anonymous HIV testing. Procedures to ensure client anonymity (i.e., no indication of testing in the client's record and no recording of personal identifying information on laboratory requests) should be developed. Even when staff members providing CTR services know the client (including name and locating information) from other activities, the client's right to be tested anonymously should be protected.

Staff Safety

Providing services in outreach settings (e.g., bars, parks) might compromise staff safety, which must be considered in development of outreach protocols. Appropriate training and precautions (e.g., working in teams) should be developed in planning services in nontraditional settings.

QUALITY ASSURANCE AND EVALUATION OF HIV CTR SERVICES

Quality Assurance

Written quality assurance protocols should be developed, made available to all staff members providing CTR services, and routinely implemented. All staff members should receive training and orientation regarding quality assurance. For information specific to ensuring high-quality CTR services, see Ensuring High-Quality HIV Prevention Counseling, Ensuring High-Quality Testing, and Ensuring High-Quality Referral Services. Quality assurance activities should address the following:

- Accessibility of services (e.g., hours of operation, location, availability of supplies and materials such as brochures, posters, test kits, safe injection materials, condoms, or lubricant).
- Compliance with written protocols for provision of service to an individual client (e.g., appropriate counseling protocols, timely return of HIV test results, referral for services responsive to client's priority needs).
- Services and materials appropriate to the client's culture, language, sex, sexual orientation, age, and developmental level.
- Staff performance/proficiency (e.g., competence, skills, credentials, and training).
- Supervision of staff members, including routine, timely feedback.
- Compliance with program guidelines and performance standards.
- Appropriateness of services to client needs, measured with client satisfaction tools (e.g., surveys or suggestion boxes).
- Record-keeping procedures, including confidentiality and security.
- Community resources (availability and collaborative arrangements).
- Collection, handling, and storage of specimens.
- Assurance of adequate funding and institutional support for CTR services.

Evaluation

CTR services should be continually evaluated to improve services to clients and provide accountability to stakeholders (150, 151). Evaluation should be interactive, involving individual persons and organizations affected by the services (150). In public health settings, the community goals outlined in community health planning processes and other relevant local planning processes could be incorporated.

Providers should identify the key, relevant program goals and objectives that reflect services to the program, community, and client, and then determine what data are needed to evaluate those goals and objectives. Information obtained from the evaluation should be used to plan and prioritize provision of CTR services within a setting. For example, information from the HIV Counseling and Testing System (133) or locally available

sources could be used during local community planning (e.g., HIV prevention community planning) to help develop or revise an HIV/AIDS prevention plan or describe who needs services. If resources for evaluation are limited, comprehensive evaluations (e.g., examining outcome or impact) might not be possible. However, even with limited resources, providers can conduct meaningful evaluations by focusing on relevant local outcomes (82).

Data

Data collected should have a clear, anticipated use and should not be the focus of or interfere with provision of CTR services. Data should be used to evaluate the extent to which the goals of CTR and locally defined service outcomes (e.g., targeted return rates, knowledge of HIV infection status, proportion of successful referrals) are met. Although sound data are essential for evaluation of services, the primary purpose of each visit should be to provide the best possible service to the client. Data should be recorded outside the time reserved for CTR discussions between the provider and the client. Clients could complete a questionnaire or intake information form on admission, providers could complete the forms immediately after meeting with a client, or a combination of the two approaches could be used.

Data collection methods should be compatible with the evaluation needs and priorities of the CTR setting and locally defined service outcomes. Data should be collected with a standard collection instrument throughout the program. Simple data collection instruments (e.g., intake forms, medical record reviews) should be developed so data can be collected unobtrusively as part of the provision of services.

Publicly funded CTR sites collect data on client demographic characteristics, risk behavior/exposure category, test acceptance, and type of site where service is provided (133). Most sites record the date of visit, anonymous or confidential test status, previous test result, current test result, and return for current test result for each client encounter. Additional data can be useful for evaluation of services, including date of previous test, type of current test (e.g., standard, rapid, oral fluid), risk-reduction plan summary, information relevant to any referrals made (e.g., provider and service description, information and materials provided, whether an appointment was made), whether the referral was received, type of service provided, dates when services were provided, and other relevant information (e.g., follow-up required, additional service needs).

Confidentiality

Any data collected or recorded must be collected or recorded in a manner that ensures the confidentiality of the client. Clear procedures and protocol manuals must be developed and used.

Ensuring High-Quality Evaluation

- The system used to collect the information must be monitored periodically to ensure data quality, which depends on the cooperative efforts of all persons providing CTR services. Periodically, data collection systems should check records at each level of the data-collection process to ensure that information is recorded consistently and completely.

- Adequate training in the use of data collection instruments should be provided to all staff members to ensure that the evaluation process is not interfering with the provision of high-quality CTR services.
- The information assembled during the evaluation process should be analyzed and reported in a timely manner to individual persons and organizations affected by the service.
- Information and feedback gained during the evaluation process should be used to improve the services offered by the site to the client.

CONCLUSION

Advances in HIV prevention and medical treatment increase the importance of HIV CTR services. Prevention counseling and knowledge of HIV status can help persons who are HIV-infected or at increased risk for HIV infection reduce their risk for transmitting or acquiring HIV infection. Referral can help persons access relevant medical, preventive, and psychosocial support services to reduce their risk for transmitting or acquiring HIV infection. These guidelines recommend how CTR can be provided to clients who could most benefit from these services across various settings and client populations.

ADDITIONAL RESOURCES

Additional information on HIV CTR can be obtained from the following sources:

- CDC's National Center for HIV, STD, and TB Prevention website at <<http://www.cdc.gov/nchstp/od/nchstp.html>>.
- CDC National AIDS Hotline in English, (800) 342-2437.
- CDC National AIDS Hotline in Spanish, (800) 344-7432.
- CDC National AIDS Hotline TTY, (800) 243-7889.
- CDC National STD Hotline, (800) 227-8922.
- CDC's National Prevention Information Network at <<http://www.cdcnpin.org>> or (800) 458-5231 (information available in English and Spanish).
- HIV/AIDS Treatment Information Service at <<http://www.hivatis.org>> or (800) 448-0440 (information available in English and Spanish).
- AIDS Clinical Trials Information Service at <<http://www.actis.org>> or (800) 874-2572 (information available in English and Spanish).
- National Clinicians' Post-Exposure Prophylaxis Hotline at <<http://pepline.ucsf.edu/PEPLine>> or (888) 448-4911.

Acknowledgments

We are grateful for the contributions of health professionals within CDC and elsewhere (e.g., Health Resources and Services Administration, National Alliance of State and Territorial AIDS Directors, academic institutions, health departments, health-care organizations). We also thank graduate students/research assistants Jennifer Chapman, M.P.H., Steven J. Connor, M.P.H., and Parag R. Sanghvi, M.S.P.H., National Center for HIV, STD, and TB Prevention, CDC.

References

1. CDC. Current trends: additional recommendations to reduce sexual and drug abuse-related transmission of human T-lymphotropic virus type III/lymphadenopathy-associated virus. *MMWR* 1986;35:152-5.
2. CDC. Perspectives in disease prevention and health promotion: Public Health Service guidelines for counseling and antibody testing to prevent HIV infection and AIDS. *MMWR* 1987;36:509-15.
3. CDC. Technical guidance on HIV counseling. *MMWR* 1993;42(No. RR-2):5-9.
4. CDC. HIV counseling, testing and referral standards and guidelines. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC, 1994.
5. Kamb ML, Fishbein M, Douglas JM Jr, et al. Efficacy of risk-reduction counseling to prevent human immunodeficiency virus and sexually transmitted diseases: a randomized controlled trial. *JAMA* 1998;280:1161-7.
6. The National Institutes of Mental Health (NIMH) Multisite HIV Prevention Trial Group. The NIMH multisite HIV prevention trial: reducing HIV sexual risk behavior. *Science* 1998;280:1889-94.
7. Boyer CB, Barrett DC, Peterman TA, Bolan G. Sexually transmitted disease (STD) and HIV risk in heterosexual adults attending a public STD clinic: evaluation of a randomized controlled behavioral risk-reduction intervention trial. *AIDS* 1997;11:359-67.
8. St. Lawrence JS, Brasfield TL, Jefferson KW, Alleyne E, O'Bannon RE, III, Shirley A. Cognitive-behavioral intervention to reduce African American adolescents' risk for HIV infection. *J Consult Clin Psychol* 1995;63:221-37.
9. McCusker J, Willis G, McDonald M, Sereti SM, Lewis BF, Sullivan JL. Community-wide HIV counselling and testing in central Massachusetts: who is retested and does their behavior change? *J Community Health* 1996;21:11-22.
10. Kelly JA, Murphy DA, Washington CD, et al. The effects of HIV/AIDS intervention groups for high-risk women in urban clinics. *Am J Pub Health* 1994;84:1918-22.
11. Tudiver F, Myers T, Kurtz RG, et al. The talking sex project: results of a randomized controlled trial of small-group AIDS education for 612 gay and bisexual men. *Evaluation and the Health Professions* 1992;15:26-42.
12. Wenger NS, Linn LS, Epstein M, Shapiro MF. Reduction of high-risk sexual behavior among heterosexuals undergoing HIV antibody testing: a randomized clinical trial. *Am J Pub Health* 1991;81:1580-5.
13. Orr DP, Langefeld CD, Katz BP, Caine VA. Behavioral intervention to increase condom use among high-risk female adolescents. *J Pediatr* 1996;128:288-95.
14. Hobfoll SE, Jackson AP, Lavin J, Britton PJ, Shepherd JB. Reducing inner-city women's AIDS risk activities: a study of single, pregnant women. *Health Psychol* 1994;13:397-403.
15. Kelly JA, St. Lawrence JS, Hood HV, Brasfield TL. Behavioral intervention to reduce AIDS risk activities. *J Consult Clin Psychol* 1989;57:60-7.
16. Valdiserri RO, Lyter DW, Leviton LC, Callahan CM, Kingsley LA, Rinaldo CR. AIDS prevention in homosexual and bisexual men: results of a randomized trial evaluating two risk reduction interventions. *AIDS* 1989;3:21-6.
17. Jemmott JB, III, Jemmott LS, Fong GT. Reductions in HIV risk-associated sexual behaviors among black male adolescents: effects of an AIDS prevention intervention. *Am J Pub Health* 1992;82:372-7.
18. Shain RN, Piper JM, Newton ER, et al. A randomized, controlled trial of a behavioral intervention to prevent sexually transmitted disease among minority women. *N Eng J Med* 1999;340:93-100.
19. O'Donnell C, O'Donnell L, San Doval A, Duran R, Labes K. Reductions in STD infections subsequent to an STD clinic visit: using video-based patient education to supplement provider interactions. *Sex Transm Dis* 1998;25:161-7.

20. Cohen DA, MacKinnon DP, Dent C, Mason HRC, Sullivan E. Group counseling at STD clinics to promote use of condoms. *Public Health Rep* 1992;107:727–31.
21. Cohen D, Dent C, MacKinnon D. Condom skills education and sexually transmitted disease reinfection. *J Sex Research* 1991;28:139–44.
22. Palella FJ Jr, Delaney KM, Moorman AC, et al. Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. *N Eng J Med* 1998;338:853–60.
23. CDC. 1999 USPHS/IDSA guidelines for the prevention of opportunistic infections in persons infected with human immunodeficiency virus. *MMWR* 1999;48(No. RR-10):1–59.
24. Connor EM, Sperling RS, Gelber R, et al. Reduction of maternal-infant transmission of human immunodeficiency virus type 1 with zidovudine treatment. *N Eng J Med* 1994;331:1173–80.
25. Kassler WJ. Advances in HIV testing technology and their potential impact on prevention. *AIDS Educ Prev* 1997;9(suppl B):27–40.
26. Spielberg B, Goldbaum G, Branson B, Wood B. Acceptance of alternate HIV counseling and testing strategies [Abstract]. Presented in the 1999 National HIV Prevention Conference, Atlanta, GA, 1999.
27. CDC. HIV partner counseling and referral services: guidance. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC, 1998.
28. CDC. HIV prevention case management: guidance. Atlanta, GA: US Department of Health and Human Services, CDC, 1997.
29. CDC. 1998 guidelines for treatment of sexually transmitted diseases. *MMWR* 1998;47(No. RR-1):1–30. (These guidelines will be updated in 2002 and available at <<http://www.cdc.gov/nchstp/od/nchstp.html>>.)
30. CDC. Recommendations for prevention and control of hepatitis C virus (HCV) infection and HCV-related chronic disease. *MMWR* 1998;47(No. RR-19):1–39.
31. CDC. Prevention of hepatitis A through active or passive immunization: recommendations of the Immunization Practices Advisory Committee (ACIP). *MMWR* 1999;48(No. RR-12):1–37.
32. CDC. Hepatitis B virus: a comprehensive strategy for eliminating transmission in the United States through universal childhood vaccination: recommendations of the Immunization Practices Advisory Committee (ACIP). *MMWR* 1991;40(No. RR-13):1–25.
33. US Preventive Services Task Force. DiGuseppi C, Atkins D, Woolf SH, eds. *Guide to clinical preventive services*. 2nd ed. Baltimore, MD: Williams & Wilkins, 1996.
34. CDC. *CDC guidelines: improving the quality*. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC, 1996.
35. CDC. HIV and AIDS—United States, 1981–2000. *MMWR* 2001;50:430–4.
36. CDC. Guidelines for national human immunodeficiency virus case surveillance, including monitoring for human immunodeficiency virus infection and acquired immunodeficiency syndrome. *MMWR* 1999;48(No. RR-13):1–28.
37. Collis TK, Celum CL. The clinical manifestations and treatment of sexually transmitted diseases in human immunodeficiency virus-positive men. *Clin Infect Dis* 2001;32:611–22.
38. Valdiserri RO, Holtgrave DR, West GR. Promoting early HIV diagnosis and entry into care. *AIDS* 1999;13:2317–30.
39. Rietmeijer CA, Kane MS, Simons PZ, et al. Increasing the use of bleach and condoms among injecting drug users in Denver: outcomes of a targeted, community-level HIV prevention program. *AIDS* 1996;10:291–8.
40. Rhodes F, Malotte CK. HIV risk interventions for active drug users. In: S.Oskamp, S.Thompson, eds. *Understanding HIV risk behavior: safer sex and drug use*. Thousand Oaks, CA: Sage Publications, 1996:297–36.
41. Gibson DR, Lovelle-Drache J, Young M, Hudes ES, Sorensen JL. Effectiveness of brief counseling in reducing HIV risk behavior in injecting drug users: final results of randomized trials of counseling with and without HIV testing. *AIDS and Behavior* 1999;3:3–12.

42. Doll LS, O'Malley PM, Pershing AL, Darrow WW, Hessel NA, Lifson AR. High-risk sexual behavior and knowledge of HIV antibody status in the San Francisco City Clinic Cohort. *Health Psychol* 1990;9:253-65.
43. Cleary PD, Van Devanter N, Rogers TF, et al. Behavior changes after notification of HIV infection. *Am J Pub Health* 1991;81:1586-90.
44. Fox R, Odaka NJ, Brookmeyer R, Polk BF. Effect of HIV antibody disclosure on subsequent sexual activity in homosexual men. *AIDS* 1987;1:241-6.
45. van Griensven GJP, de Vroome EMM, Tielman RAP, et al. Effect of human immunodeficiency virus (HIV) antibody knowledge on high-risk sexual behavior with steady and nonsteady sexual partners among homosexual men. *Am J Epidemiol* 1989;129:596-603.
46. Coates TJ, Morin SF, McKusick L. Behavioral consequences of AIDS antibody testing among gay men [Letter]. *JAMA* 1987;258:1889.
47. Wenger NS, Kusseling FS, Beck K, Shapiro MF. Sexual behavior of individuals infected with the human immunodeficiency virus: the need for intervention. *Arch Intern Med* 1994;154:1849-54.
48. Desenclos J-C, Papaevangelou G, Ancelle-Park R, for the European Community Study Group on HIV in Injecting Drug Users. Knowledge of HIV serostatus and preventive behaviour among European injecting drug users. *AIDS* 1993;7:1371-7.
49. Dawson J, Fitzpatrick R, McLean J, Hart G, Boulton M. The HIV test and sexual behavior in a sample of homosexually active men. *Soc Sci Med* 1991;32:683-8.
50. Otten MW Jr, Zaidi AA, Wroten JE, Witte J, Peterman TA. Changes in sexually transmitted disease rates after HIV testing and posttest counseling, Miami, 1988 to 1989. *Am J Pub Health* 1993;83:529-33.
51. Quinn TC, Wawer MJ, Sewankambo N, et al. Viral load and heterosexual transmission of human immunodeficiency virus type 1. *N Eng J Med* 2000;342:921-9.
52. Branson BM. Home sample collection tests for HIV infection. *JAMA* 1998;280:1699-701.
53. CDC. Prevention and treatment of tuberculosis among patients infected with human immunodeficiency virus: principles of therapy and revised recommendations. *MMWR* 1998;47(No. RR-20):1-25.
54. CDC. Updated U.S. Public Health Service guidelines for the management of occupational exposures to HBV, HCV, and HIV and recommendations for postexposure prophylaxis. *MMWR* 2001;50(No. RR-11):1-54.
55. CDC. Management of possible sexual, injecting-drug-use, or other nonoccupational exposure to HIV, including considerations related to antiretroviral therapy. *MMWR* 1998;47(No. RR-17):1-14.
56. Tao G, Irwin KL, Kassler WJ. Missed opportunities to assess STDs in US adults during routine medical checkups. *Am J Prev Med* 2000;18:109-14.
57. CDC. HIV prevention practices of primary care physicians—United States, 1992. *MMWR* 1994;42:988-92.
58. Tiara DA, Safran DG, Seto TB, Rogers WH, Tarlov AR. The relationship between patient income and physician discussion of health risk behaviors. *JAMA* 1997;278:1412-7.
59. Schwartz JS, Lewis CE, Clancy C, Kinosian MS, Radany MH, Koplan JP. Internists' practices in health promotion and disease prevention. *Ann Intern Med* 1991;114:46-53.
60. Bindman AB, Osmond D, Hecht FM, et al. Multistate Evaluation of anonymous HIV testing and access to medical care. *JAMA* 1998;280:1416-20.
61. Chen Z, Branson B, Ballenger A, Peterman TA. Risk assessment to improve targeting of HIV counseling and testing services of STD clinic patients. *Sex Transm Dis* 1998;25:539-43.
62. Quinn TC, Glasser D, Cannon RO, et al. Human immunodeficiency virus infection among patients attending clinics for sexually transmitted diseases. *N Eng J Med* 1988;318:197-203.

63. Erickson B, Wasserheit JN, Rompalo AM, Brathwaite W, Glasser D, Hook EW III. Routine voluntary HIV screening in STD clinic clients: characterization of infected clients. *Sex Transm Dis* 1990;17:194-9.
64. Groseclose S, Erickson B, Quinn T, Glasser D, Campbell C, Hook E. Characteristics of patients accepting and refusing routine, voluntary HIV antibody testing in public sexually transmitted disease clinics. *Sex Transm Dis* 1994;21:31-5.
65. Kassler WJ, Zenilman JM, Erickson B, Fox R, Peterman TA, Hook EW III. Seroconversion in patients attending sexually transmitted disease clinics. *AIDS* 1994;8:351-5.
66. Asch SM, London AS, Barnes PF, Gelberg L. Testing for human immunodeficiency virus infection among tuberculosis patients in Los Angeles. *Am J Respir Crit Care Med* 1997;155:378-81.
67. Pitchenik AE, Burr J, Suarez M, Fertel D, Gonzalez G, Moas C. Human T-cell lymphotropic virus-III (HTLV-III) seropositivity and related disease among 71 consecutive patients in whom tuberculosis was diagnosed. *Am Rev Respir Dis* 1987;135:875-9.
68. Shafer RW, Chirgwin KD, Glatt AE, Dahdouh MA, Landesman SH, Suster B. HIV prevalence, immunosuppression, and drug resistance in patients with tuberculosis in an area endemic for AIDS. *AIDS* 1991;5:399-405.
69. Theuer CP, Hopewell PC, Elias D, Schechter GF, Rutherford GW, Chaisson RE. Human immunodeficiency virus infection in tuberculosis patients. *J Infect Dis* 1990;162:8-12.
70. Peterman TA, Todd KA, Mupanduki I. Opportunities for targeting publicly funded human immunodeficiency virus counseling and testing. *J Acquir Immune Defic Syndr* 1996;12:69-74.
71. Janssen RS, St. Louis ME, Satten GA, et al. HIV infection among patients in U.S. acute care hospitals: strategies for the counseling and testing of hospital patients. *New Eng J Med* 1992;327:445-52.
72. CDC. HIV prevalence trends in selected populations in the United States: results from national serosurveillance, 1993-1997. Atlanta, GA: US Department of Health and Human Services, CDC, 2001.
73. Kelly JA, St. Lawrence JS. The prevention of AIDS: roles for behavioral intervention. *Scand J Behav Therapy* 1987;16:5-19.
74. Cohen DA, Dent C, MacKinnon D, Hahn G. Condoms for men, not women: results of brief promotion programs. *Sex Transm Dis* 1992;19:245-51.
75. Sikkema KJ, Bissett RT. Concepts, goals, and techniques of counseling: review and implications for HIV counseling and testing. *AIDS Educ Prev* 1997;9(suppl B):14-26.
76. Roffman RA, Kalichman SC, Kelly JA, et al. HIV antibody testing of gay men in smaller US cities. *AIDS Care* 1995;7:405-13.
77. DiClemente RJ, Wingood GM. A randomized controlled trial of an HIV sexual risk-reduction intervention for young African-American women. *JAMA* 1995;274:1271-6.
78. Kelly JA, St. Lawrence JS, Diaz YE, et al. HIV risk behavior reduction following intervention with key opinion leaders of population: an experimental analysis. *Am J Pub Health* 1991;81:168-71.
79. Kelly JA, St. Lawrence JS. Behavioral intervention and AIDS. *The Behavioral Therapist* 1986;6:121-5.
80. Kamb ML, Dillon BA, Fishbein M, Willis KL, and the Project RESPECT Study Group. Quality assurance of HIV prevention counseling in a multi-center randomized controlled trial. *Public Health Rep* 1996;111(suppl 1):99-107.
81. Holtgrave DR, Valdiserri RO, Gerber AR, Hinman AR. Human immunodeficiency virus counseling, testing, referral, and partner notification services: a cost-benefit analysis. *Arch Intern Med* 1993;153:1225-30.
82. Holtgrave DR, Reiser WJ, DiFranceisco W. The evaluation of HIV counseling-and-testing services: making the most of limited resources. *AIDS Educ Prev* 1997;9(3 suppl):105-18.

83. Kamb ML, Kassler W, Peterman TA, and the Project RESPECT Study Group. Cost of preventing HIV via counseling: results from a randomized trial (Project RESPECT) [Abstract 33263]. Presented at the XII International Conference on AIDS, Geneva, Switzerland, 1998:644.
84. Booth RE, Kwiatkowski CF, Stephens RC. Effectiveness of HIV/AIDS interventions on drug use and needle risk behaviors for out-of-treatment injection drug users. *J Psychoactive Drugs* 1998;30:269–78.
85. Castrucci BC, Kamb ML, Hunt K. Assessing use of the 1994 HIV counseling, testing, and referral standards and guidelines—how closely does practice conform to existing recommendations? [Abstract P125]. Presented at the 2000 National STD Prevention Conference, December 4–7, Milwaukee, WI, 2000.
86. Kelly JA, Murphy DA, Sikkema KJ, Kalichman SC. Psychological interventions to prevent HIV infection are urgently needed: new priorities for behavioral research in the second decade of AIDS. *Am Psychol* 1993;48:1023–34.
87. McCusker J, Stoddard AM, Zapka JG, Zorn M, Mayer KH. Predictors of AIDS-preventive behavior among homosexually active men: a longitudinal study. *AIDS* 1989;3:443–8.
88. Kelly JA, Murphy DA. Some lessons learned about risk reduction after ten years of the HIV/AIDS epidemic. *AIDS Care* 1991;3:251–7.
89. Kelly JA, Murphy DA. Psychological interventions with AIDS and HIV: prevention and treatment. *J Consult Clin Psychol* 1992;60:576–85.
90. American Public Health Association. AIDS prevention in the community: lessons from the first decade. Washington, DC: American Public Health Association, 1995.
91. Wiktor SZ, Biggar RJ, Melbye M, et al. Effect of knowledge of human immunodeficiency virus infection status on sexual activity among homosexual men. *J Acquir Immune Defic Syndr* 1990;3:62–8.
92. Kelly JA, St. Lawrence JS, Betts R, Brasfield TL, Hood HV. A skills-training group intervention model to assist persons in reducing risk behaviors for HIV infection. *AIDS Educ Prev* 1990;2:24–35.
93. Sikkema KJ, Winett RA, Lombard DN. Development and evaluation of an HIV-risk reduction program for female college students. *AIDS Educ Prev* 1995;7:145–59.
94. Kelly JA, Kalichman SC. Increased attention to human sexuality can improve HIV-AIDS prevention efforts: key research issues and directions. *J Consult Clin Psychol* 1995;63:907–18.
95. CDC. Compendium of HIV prevention interventions with evidence of effectiveness. Atlanta, GA: US Department of Health and Human Services, CDC, 1999.
96. Higgins DL, Galavotti C, O'Reilly KR, et al. Evidence for the effects of HIV antibody counseling and testing on risk behaviors. *JAMA* 1991;266:2419–29.
97. Wolitski RJ, MacGowan RJ, Higgins DL, Jorgenson CM. The effects of HIV counseling and testing on risk-related practices and help-seeking behavior. *AIDS Educ Prev* 1997;9(suppl B):52–67.
98. Flaskerud JH. Matching client and therapist ethnicity, language, and gender: a review of research. *Issues in Mental Health Nursing* 1990;11:321–36.
99. Kilmarx PH, Hamers FF, Peterman TA. Living with HIV: experiences and perspectives of HIV-infected sexually transmitted disease clinic patients after posttest counseling. *Sex Transm Dis* 1998;25:28–37.
100. Cates W, Handsfield HH. HIV counseling and testing: does it work? *Am J Public Health* 1988;78:1533–4.
101. Calsyn DA, Saxon AJ, Freeman G, Whittaker S. Ineffectiveness of AIDS education and HIV antibody testing in reducing high risk behaviors among injection drug users. *Am J Public Health* 1992;82:573–5.
102. Edlin BR, Irwin KL, Faruque S, et al. Intersecting epidemics—crack cocaine use and HIV infection among inner-city young adults. *N Eng J Med* 1994;331:1422–7.

103. Cottler LB, Leukefeld C, Hoffman J, et al. Effectiveness of HIV risk reduction initiatives among out-of-treatment non-injection drug users. *J Psychoactive Drugs* 1998;30:279-90.
104. Nicolosi A, Molinari S, Musicco M, Saracco A, Ziliani N, Lazzarin A. Positive modification of injecting behavior among intravenous heroin users from Milan and Northern Italy 1987-1989. *Brit J Addiction* 1991;86:91-102.
105. Neaigus A, Sufian M, Friedman S, et al. Effects of outreach intervention on risk reduction among intravenous drug users. *AIDS Educ Prev* 1990;2:253-71.
106. Obermeyer TE, Streeter A. Street outreach HIV education to intravenous drug users and other substance abusers. *AIDS Patient Care* 1991;5:312-4.
107. Hagan H, Jarlais DC, Friedman SR, Purchase D, Alter MJ. Reduced risk of hepatitis B and hepatitis C among injection drug users in the Tacoma Syringe Exchange Program. *Am J Public Health* 1995;85:1531-7.
108. Jones TS, Vlahov D. Use of sterile syringes and aseptic drug preparation are important components of HIV prevention among injection drug users. *J Acquir Immune Defic Syndr* 1998;18(suppl 1):S1-S5.
109. Padian NS, O'Brien TR, Chang Y, Glass S, Francis DP. Prevention of heterosexual transmission of human immunodeficiency virus through couple counseling. *J Acquir Immune Defic Syndr* 1993;6:1043-8.
110. Kamenga M, Ryder RW, Jingu M, et al. Evidence of marked sexual behavior change associated with low HIV-1 seroconversion in 149 married couples with discordant HIV-1 serostatus: experience at an HIV counselling center in Zaire. *AIDS* 1991;5:61-7.
111. De Vincenzi I. A longitudinal study of human immunodeficiency virus transmission by heterosexual partners. *N Eng J Med* 1994;331:341-6.
112. Levy J, Fox S, Valle M. What you don't know can hurt you: the influence of prior HIV testing on serostatus results at repeat testing [Abstract 43113]. Presented at the 12th World AIDS Conference, Geneva, Switzerland, 1998:869.
113. Irwin KL, Valdiserri RO, Holmberg SD. The acceptability of voluntary HIV antibody testing in the United States: a decade of lessons learned. *AIDS* 1996;10:1707-17.
114. Valdiserri RO, Moore M, Gerber AR, Campbell CH Jr, Dillon BA, West GR. A study of clients returning for counseling after HIV testing: implications for improving rates of return. *Public Health Rep* 1993;108:12-8.
115. Catania JA, Gibson DR, Marin B, Coates TJ, Greenblatt RM. Response bias in assessing sexual behaviors relevant to HIV transmission. *Evaluation and Program Planning* 1990;13:19-29.
116. Weber JT, Frey R Jr, Horsley R, Gwinn ML. Publicly funded HIV counseling and testing in the United States, 1992-1995. *AIDS Educ Prev* 1997;9(suppl B):79-91.
117. Branson B, Ballenger A, Olthoff G. HIV test results and post-test counseling by telephone [Abstract PC0535]. Presented at the Tenth International Conference on AIDS, 1994.
118. Schluter WW, Judson FN, Baron AE, McGill WL, Marine WM, Douglas JM Jr. Usefulness of human immunodeficiency virus post-test counseling by telephone for low-risk clients of an urban sexually transmitted diseases clinic. *Sex Transm Dis* 1996;23:190-7.
119. Samet JH, Freedberg KA, Stein MD, et al. Trillion virion delay: time from testing positive for HIV to prevention for primary care. *Arch Intern Med* 1998;158:734-40.
120. CDC. Update: HIV counseling and testing using rapid tests—United States, 1995. *MMWR* 1998;47:211-5.
121. Federal Trade Commission. Home-use tests for HIV can be inaccurate, FTC warns [Consumer Alert]. 1999. Available at <<http://www.ftc.gov/bcp/conline/pubs/alerts/hivalrt.htm>>. Accessed July 13, 2001.
122. CDC. Identification of HIV-1 group O infection—Los Angeles County, California, 1996. *MMWR* 1996;45:561-5.
123. Sullivan PS, Do AN, Robbins K, et al. Surveillance for variant strains of HIV: subtype G and group O HIV-1 [Letter]. *JAMA* 1997;278:292.

124. CDC. Human immunodeficiency virus type 2. Atlanta, GA: US Department of Health and Human Services, CDC, 1998. Available at <<http://www.cdc.gov/hiv/pubs/facts/hiv2.htm>>. Accessed July 4, 2001.
125. CDC. Testing for antibodies to human immunodeficiency virus type 2 in the United States. *MMWR* 1992;41(No. RR-12):1-9.
126. CDC. U.S. Public Health Service guidelines for testing and counseling of blood and plasma donors for human immunodeficiency virus type 1 antigen. *MMWR* 1996;45(No. RR-2):1-9.
127. Celum CL, Coombs RW, Lafferty W, et al. Indeterminate human immunodeficiency virus type 1 Western blots: seroconversion risk, specificity of supplemental tests, and an algorithm for evaluation. *J Infect Dis* 1991;164:656-64.
128. Stetler HC, Granada TC, Nunez CA, et al. Field evaluation of rapid HIV serologic tests for screening and confirming HIV-1 infection in Honduras. *AIDS* 1997;11:369-75.
129. Horsburgh CR Jr, Jason J, Longini I, et al. Duration of human immunodeficiency virus infection before detection of antibody. *Lancet* 1989;2:637-40.
130. Busch MP, Lee LLJ, Satten GA, et al. Time course of detection viral and serologic markers preceding human immunodeficiency virus type 1 seroconversion: implications for screening of blood and tissue donors. *Transfusion* 1995;35:91-7.
131. Jackson JB, MacDonald KL, Cadwell J, et al. Absence of HIV infection in blood donors with indeterminate Western blot tests for antibody to HIV-1. *N Eng J Med* 1990;322:217-22.
132. Dock NL, Kleinman SH, Rayfield MA, Schable CA, Williams AE, Dodd RY. Human immunodeficiency virus infection and indeterminate Western blot patterns: prospective studies in a low prevalence population. *Arch Intern Med* 1991;151:525-30.
133. CDC. HIV counseling and testing in publicly funded sites: annual report, 1997 and 1998. Atlanta, GA: US Department of Health and Human Services, CDC, 2001.
134. Ciesielski CA, Metler RP. Duration of time between exposure and seroconversion in healthcare workers with occupationally acquired infection with human immunodeficiency virus. *Am J Med* 1997;102:115-6.
135. CDC. HIV testing among populations at risk for HIV infection—nine states, November 1995–December 1996. *MMWR* 1998;47:1086-91.
136. Wykoff RF, Jones JL, Longshore ST, et al. Notification of the sex and needle-sharing partners of individuals with human immunodeficiency virus in rural South Carolina: 30 month experience. *Sex Transm Dis* 1991;18:217-22.
137. Exner TM, Ehrhardt A, Loeb I, Zawadzki R. HIV counseling and testing: women's experiences and the role of testing as a prevention strategy [Abstract We.C.3529]. In: *Proceedings of the XI International Conference on AIDS, 1996;11(2):150*.
138. Kassler WJ, Meriwether RA, Klimko TB, Peterman TA, Zaidi A. Eliminating access to anonymous HIV antibody testing in North Carolina: effects on HIV testing and partner notification. *J Acquir Immune Defic Syndr* 1997;14:281-9.
139. Simpson WM, Johnstone FD, Goldberg DJ, Gormley SM, Hart GJ. Antenatal HIV testing: assessment of a routine voluntary approach. *BMJ* 1999;318:1660-1.
140. Lee JH, Mitchell B, Nolt B, Robbins B, Thomas MC, Branson BM. Targeted opt-in vs. routine opt-out HIV testing in an STD clinic [Abstract 153]. Presented at the 1999 National HIV Prevention Conference, August 29–September 1, Atlanta, GA, 1999.
141. Rahimian A, Driscoll M, Taylor D, Cohen M. Barriers to building a comprehensive system of HIV counseling and testing by consent to women of reproductive age in Chicago, Illinois [Abstract Tu.D. 2772]. In: *Proceedings of the XI International Conference on AIDS, 1996;1:396*.
142. Nutting PA, Main DS, Fischer PM, et al. Problems in laboratory testing in primary care. *JAMA* 1996;275:635-9.

143. Boone DJ, Steindel SD, Herron R, et al. Transfusion medicine monitoring practices: a study of the College of American Pathologists/CDC Outcomes Working Group. *Arch Path Lab Med* 1995;119:999–1006.
144. Witte DL, VanNess SA, Angstandt DS, Pennell BJ. Errors, mistakes, blunders, outliers, or unacceptable results: how many. *Clin Chem* 1997;43:1352–6.
145. Schochetman G, George JR, eds. *AIDS testing: a comprehensive guide to technical, medical, social, legal, and management issues*. 2 ed. New York, NY: Springer-Verlag, 1994.
146. Institute of Medicine. Eng TR, Butler WT, eds. *The hidden epidemic: confronting sexually transmitted diseases*. Washington, DC: National Academy Press, 1997.
147. Greenberg JB, MacGowan R, Neumann M, et al. Linking injection drug users to medical services: role of street outreach referrals. *Health Soc Work* 1998;23:298–309.
148. Marx R, Hirozawa AM, Chu PL, Bolan GA, Katz M. Linking clients for HIV antibody counseling and testing to prevention services. *J Community Health* 1999;24:201–14.
149. Hymel MS, Greenberg BL. The Walden House Young Adult HIV Project: meeting the needs of multidagnosed youth. *J Adolesc Health* 1998;23S:122–31.
150. CDC. Framework for program evaluation in public health. *MMWR* 1999;48(No. RR-11):1–40.
151. CDC. *Evaluating CDC-funded health department HIV prevention programs. Volume 1: guidance*. Atlanta, GA: US Department of Health and Human Services, CDC, 1999.

Glossary

AIDS: Acquired immunodeficiency syndrome. AIDS can affect the immune and central nervous systems and can result in neurological problems, infections, or cancers. It is caused by human immunodeficiency virus (HIV).

Anal sex: A type of sexual intercourse in which a man inserts his penis in his partner's anus. Anal sex can be insertive or receptive.

Anonymous: In anonymous testing, client identifying information is not linked to testing information, including the request for tests or test results.

Antiretroviral therapy: Treatment with drugs designed to prevent HIV from replicating in HIV-infected persons. Highly active antiretroviral therapy (HAART) is an antiretroviral regimen that includes multiple classifications of antiretroviral drugs.

Client-centered HIV prevention counseling: An interactive risk-reduction counseling model usually conducted with HIV testing, in which the counselor helps the client identify and acknowledge personal HIV risk behaviors and commit to a single, achievable behavior change step that could reduce the client's HIV risk.

Confidentiality: Pertains to the disclosure of personal information in a relationship of trust and with the expectation that it will not be divulged to others in ways that are inconsistent with the original disclosure. Confidentiality must be maintained for persons who are recommended and/or who receive HIV counseling, testing, and referral (CTR) services.

Confidential HIV test: An HIV test for which a record of the test and the test results are recorded in the client's chart.

Confirmatory test: A highly specific test designed to confirm the results of an earlier (screening) test. For HIV testing, a Western blot or, less commonly, an immunofluorescence assay (IFA) is used as a confirmatory test.

EIA: Enzyme immunoassay. Sometimes referred to as ELISA (see next definition). A commonly used screening test to detect antibodies to HIV.

ELISA: Enzyme-linked immunosorbent assay. A type of EIA (see previous definition). A commonly used screening test to detect antibodies to HIV.

Evaluation: A process for determining how well health systems, either public or private, deliver or improve services and for demonstrating the results of resource investments.

False negative: A negative test result for a person who is actually infected.

False positive: A positive test result for a person who is actually not infected.

Freestanding HIV test site: A site that provides only HIV services. Sometimes referred to as alternate test site or anonymous test site.

HIV: Human immunodeficiency virus, which causes AIDS. Several types of HIV exist, with HIV-1 being the most common in the United States.

HIV test: More correctly referred to as an HIV antibody test, the HIV test is a laboratory procedure that detects antibodies to HIV, rather than the virus itself.

HIV prevention counseling: An interactive process between client and counselor aimed at reducing risky sex and needle-sharing behaviors related to HIV acquisition (for HIV-uninfected clients) or transmission (for HIV-infected clients). See also client-centered HIV prevention counseling.

Home sample collection test: A test that a consumer purchases and uses to collect blood (or other bodily fluid) and then send it out for testing. Counseling and test results are typically provided by telephone using user-generated codes to ensure confidentiality and anonymity.

Incidence: In epidemiology, the number of new cases of infection or disease that occur in a defined population within a specified time.

Indeterminate test result: A possible result of a Western blot, which might represent a recent HIV infection or a false-positive.

Information: In the context of HIV counseling, information encompasses the topics HIV transmission and prevention and the meaning of HIV test results.

Informed consent: The legally effective permission of a client or legally authorized representative (e.g., parent or legal guardian of a minor child) to undergo a medical test or procedure.

Negative predictive value: A negative predictive value estimates the probability that a person with a negative diagnostic test result will actually not be infected.

Nonoccupational HIV exposure: A reported sexual, injection-drug-use, or other non-occupational HIV exposure that might put a patient at high risk for acquiring HIV infection.

Nucleic acid amplification testing: A type of testing that identifies viral genes (e.g., specific sequences of nucleic acids) using gene amplification technologies such as polymerase chain reaction (PCR).

Occupational HIV exposure: An occupational exposure to HIV that occurs during the performance of job duties. Defined as a percutaneous injury (e.g., a needlestick or cut with a sharp object), contact of mucous membranes, or contact of skin (especially when the exposed skin is chapped, abraded, or afflicted with dermatitis or the contact is prolonged or involving an extensive area) with blood, tissues, or other body fluids to which universal precautions apply.

Oral fluid test: A test using oral mucosal transudate, a serous fluid. To differentiate this fluid from saliva, an absorbent material is left in the mouth for several minutes. In an HIV-infected person, oral mucosal transudate is likely to contain HIV antibodies.

Oral sex: A type of sexual intercourse in which the partner's genitals are stimulated by mouth and tongue.

Partner counseling and referral services (PCRS): A prevention activity that aims to a) provide services to HIV-infected persons and their sex and needle-sharing partners so they can reduce their risk for infection or, if already infected, can prevent transmission to others and b) help partners gain earlier access to individualized counseling, HIV testing, medical evaluation, treatment, and other prevention and support services.

Perinatal HIV transmission: Transmission of HIV from the mother to the fetus or infant during pregnancy, delivery, or breast-feeding.

Positive predictive value: A positive predictive value estimates the probability that a person with a positive diagnostic test result will actually be infected.

Positive test: For HIV, a specimen sample that is reactive on an initial ELISA test, repeatedly reactive on a second ELISA run on the same specimen, and confirmed positive on Western blot or other supplemental test indicates that the client is infected.

Prevalence: The number or percentage of persons in a given population with a disease or condition at a given point in time.

Prevention case management (PCM): A client-centered HIV prevention activity that promotes adoption of HIV risk-reduction behaviors by clients with multiple, complex problems and risk-reduction needs. PCM is a hybrid of HIV prevention counseling and traditional case management that provides intensive, on-going, individualized prevention counseling, support, and referral to other needed services.

Prevention counseling: An interactive process between client and counselor aimed at reducing risky sex and needle-sharing behaviors related to HIV acquisition (for HIV-uninfected clients) or transmission (for HIV-infected clients). See also client-centered HIV prevention counseling and HIV prevention counseling.

Quality assurance: An ongoing process for ensuring that the CTR program effectively delivers a consistently high level of service to the clients.

Rapid HIV test: A test to detect antibodies to HIV that can be collected and processed within a short interval of time (e.g., approximately 10–60 minutes).

Referral: The process through which a client is connected with services to address prevention needs (medical, prevention, and psychosocial support).

Risk assessment: Risk assessment is a fundamental part of a client-centered HIV prevention counseling session in which the client is encouraged to identify, acknowledge, and discuss in detail his or her personal risk for acquiring or transmitting HIV.

Risk screening: A brief evaluation of HIV risk factors, both behavioral and clinical, used for decisions about who should be recommended HIV counseling and testing. Risk screening is different from risk assessment.

Screening test: An initial test, usually designed to be sensitive, to identify all persons with a given condition or infection (e.g., enzyme immunoassay [EIA] or enzyme-linked immunosorbent assay [ELISA]).

Sensitivity: The probability that a test will be positive when infection or condition is present.

Seroconversion: Initial development of detectable antibodies specific to a particular antigen; the change of a serologic test result from negative to positive as a result of antibodies induced by the introduction of antigens or microorganisms into the host.

Specificity: The probability that a test will be negative when the infection or condition is not present.

Tuberculosis (TB) disease: Active disease caused by *Mycobacterium tuberculosis*, as evidenced by a confirmatory culture, or, in the absence of culture, suggestive clinical symptoms, including productive cough lasting ≥ 3 weeks, chest pain, hemoptysis, fever, night sweats, weight loss, and easy fatigability. Active TB is a communicable disease that is treatable, curable, and preventable, and persons with active TB disease should be under the care of a health-care provider. Active TB disease could indicate immune deficiency. For HIV-infected persons, active TB disease is considered an opportunistic infection and a qualifying condition for AIDS.

Tuberculosis (TB) infection: Infection with the bacteria *M. tuberculosis*, as evidenced by a positive tuberculin skin test (TST) that screens for infection with this organism. Sometimes, TST is called a purified protein derivative (PPD) or Mantoux test. A positive skin test might or might not indicate active TB disease (see tuberculosis disease). Thus, any person with a positive TST should be screened for active TB and, once active TB is excluded, evaluated for treatment to prevent the development of TB disease. TB infection alone is not considered an opportunistic infection indicating possible immune deficiency.

Vaginal sex: A type of sexual intercourse in which the man's penis enters the woman's vagina.

Voluntary HIV testing: HIV testing that is offered free of coercion. With voluntary HIV testing, participants have the opportunity to accept or refuse HIV testing.

Western blot: A laboratory test that detects specific antibodies to components of a virus. Chiefly used to confirm HIV antibodies in specimens found repeatedly reactive using ELISA.